



Branching-time logic ECTL[#] and its tree-style one-pass tableau: Extending fairness expressibility of ECTL⁺



Alexander Bolotov^{a,1}, Montserrat Hermo^{b,*,2}, Paqui Lucio^{b,2}

^a University of Westminster, W1W 6UW, London, UK

^b University of the Basque Country, P. Manuel de Lardizabal, 1, 20018-San Sebastián, Spain

ARTICLE INFO

Article history:

Received 15 February 2019

Received in revised form 23 December 2019

Accepted 8 February 2020

Available online 13 February 2020

Keywords:

Temporal logic

Fairness

Tableau

Branching-time

One-pass tableau

ABSTRACT

Temporal logic has become essential for various areas in computer science, most notably for the specification and verification of hardware and software systems. For the specification purposes rich temporal languages are required that, in particular, can express fairness constraints. For linear-time logics which deal with fairness in the linear-time setting, one-pass and two-pass tableau methods have been developed. In the repository of the CTL-type branching-time setting, the well-known logics ECTL and ECTL⁺ were developed to explicitly deal with fairness. However, due to the syntactical restrictions, these logics can only express restricted versions of fairness. The logic CTL*, often considered as 'the full branching-time logic' overcomes these restrictions on expressing fairness. However, CTL* is extremely challenging for the application of verification techniques, and the tableau technique, in particular. For example, there is no one-pass tableau construction for CTL*, while one-pass tableau has an additional benefit enabling the formulation of dual sequent calculi that are often treated as more 'natural' being more friendly for human understanding. These two considerations lead to the following problem - are there logics that have richer expressiveness than ECTL⁺, allowing the formulation of a new range of fairness constraints with 'until' operator, yet 'simpler' than CTL*, and for which a one-pass tableau can be developed? Here we give a positive answer to this question, introducing a sub-logic of CTL* called ECTL[#], its tree-style one-pass tableau, and an algorithm for obtaining a systematic tableau, for any given admissible branching-time formulae. We prove the termination, soundness and completeness of the method. As tree-shaped one-pass tableaux are well suited for the automation and are amenable for the implementation and for the formulation of sequent calculi. Our results also open a prospect of relevant developments of the automation and implementation of the tableau method for ECTL[#], and of a dual sequent calculi.

© 2020 Elsevier B.V. All rights reserved.

* Corresponding author.

E-mail addresses: A.Bolotov@westminster.ac.uk (A. Bolotov), montserrat.hermo@ehu.eu (M. Hermo), paqui.lucio@ehu.eu (P. Lucio).

¹ The author would like to thank the University of Westminster for supporting the sabbatical in 2017.

² These authors have been partially supported by Spanish Projects TIN2013-46181-C2-2-R and TIN2017-86727-C2-2-R, and by the University of the Basque Country under Project LoRea GIU15/30.

1. Introduction

Temporal logic has become essential for the specification and verification of hardware and software systems. For the specification of the reactive and distributed systems, or, most recently, autonomous systems, the modelling of the possibilities ‘branching’ into the future is essential. Among important properties of these systems, so called fairness properties are important. In the standard formalisation of fairness, operators \diamond (eventually) and \square (always) have been used: $A\diamond\square p - ‘p’$ is true along all computation paths except possibly their finite initial interval, where ‘A’ is ‘for all paths’ quantifier, and $E\square\diamond p - ‘p’$ is true along a computation path at infinitely many states, where ‘E’ stands for ‘there exists a path’ quantifier. Branching-time logics (BTL) here give us an appropriate reasoning framework, where the most used class of formalisms are ‘CTL’ (Computation Tree Logic) type logics. CTL itself requires every temporal operator to be preceded by a path quantifier, thus, cannot express fairness. ECTL (Extended CTL) [1] enables simple fairness constraints but not their Boolean combinations. ECTL⁺ [2] further extends the expressiveness of ECTL allowing Boolean combinations of temporal operators and ECTL fairness constraints (but not permitting their nesting). The logic CTL*, often considered as ‘the full branching-time logic’ overcomes these restrictions on expressing fairness. However, CTL* is extremely challenging for the application of any known technique of automated reasoning. Note that, unlike fair CTL [3] which, in tackling fairness, changes the underlying trees to those with ‘fair paths’ only, ECTL and ECTL⁺ do not impose these changes.

From another perspective, the literature on fairness constraints, even in the linear-time setting, lacks the analysis of their formulation with the \mathcal{U} (‘until’) operator. To the best of our knowledge, there are only a few research papers that raise or discuss the problem. For example, [4], introduces the logic LCTL, providing an extension of liveness constraints by the ‘until’ operator. However, LCTL belongs to ‘Fair CTL-type’ logics [5]. ‘Generalised liveness assumptions, which allow to express that the conclusion $f_2\mathcal{U}f_3$ of a liveness assumption $\square(f_1 \Rightarrow (f_2\mathcal{U}f_3))$ has to be satisfied’ are addressed in [6]. The \mathcal{U} operator in the formulation of the fairness can also be found in [7] which considers the sequential composition of processes, providing the following example - the composition of processes P_1 and P_2 ‘behaves as P_1 until its termination and then behaves as P_2 ’. Finally, [8] utilises restricted linear-time fairness constraints with \mathcal{U} in the linear-time setting. We are not aware of any other analysis of fairness constraints in branching-time setting using the \mathcal{U} operator and without restricting the underlying logic to be interpreted over the ‘fair’ paths. We bridge this gap, presenting the logic ECTL[#] (we use # to indicate some restrictions on concatenations of the modalities and their Boolean combinations). It is weaker than CTL* but extends ECTL⁺ by allowing the combinations $\square(A\mathcal{U}B)$ or $A\mathcal{U}\square B$, referred to as modalities $\square\mathcal{U}$ and $\mathcal{U}\square$. This enables the formulation of stronger fairness constraints in the branching-time setting. The fairness constraint $A(p\mathcal{U}\square q)$ reads as ‘invariant q is true along all paths of the computation except possibly their finite initial interval, where p is true’. For example, the following property specifies that whenever the user of an account is requested to change the current password, either it is changed to a fresh one, or the account is deactivated:

$$A((P_n^w \mathcal{U} \square (R_n \Rightarrow A \square (P_n^{w'} \Rightarrow w \neq w'))) \vee ((L_n \wedge P_n^w) \mathcal{U} \square ((R_n \wedge (\neg P_n^{w'} \vee w' = w)) \Rightarrow \neg L_n))) \tag{1}$$

where P_n^w ($P_n^{w'}$) stands for the account n has an associated password w (w'); L_n stands for the account n is live, and R_n means the account number n is requested to change the password, Note that formula (1) represents one of the difficult cases of ECTL[#] structures - an A-disjunctive formula, see §2.

| $\mathcal{B}(\mathcal{U}, \circ)$ (CTL) extensions | $E(\square\diamond q)$ | $E(\square\diamond q \wedge \square\diamond r)$ | $A((p\mathcal{U}\square q) \vee (s\mathcal{U}\square\neg r))$ | $A\diamond(\circ p \wedge E\circ\neg p)$ |
|------------------------------------------------------------------------------|------------------------|-------------------------------------------------|---------------------------------------------------------------|------------------------------------------|
| $\mathcal{B}(\mathcal{U}, \circ, \square\diamond)$ (ECTL) | ✓ | X | X | X |
| $\mathcal{B}^+(\mathcal{U}, \circ, \square\diamond)$ (ECTL ⁺) | ✓ | ✓ | X | X |
| $\mathcal{B}^+(\mathcal{U}, \circ, \mathcal{U}\square)$ (ECTL [#]) | ✓ | ✓ | ✓ | X |
| $\mathcal{B}^*(\mathcal{U}, \circ)$ (CTL*) | ✓ | ✓ | ✓ | ✓ |

Fig. 1. Classification of CTL-type logics and their expressiveness.

Fig. 1, which utilises another temporal operator - \circ - ‘at the next moment of time’, places our logic in the hierarchy of BTL representing their expressiveness: logics are classified by using ‘ \mathcal{B} ’ for ‘Branching’, followed by the set of only allowed modalities as parameters; \mathcal{B}^+ indicates admissible Boolean combinations of the modalities and \mathcal{B}^* reflects ‘no restrictions’ in either concatenations of the modalities or Boolean combinations between them.³ Thus, $\mathcal{B}(\mathcal{U}, \circ)$ denotes the logic CTL. In this hierarchy ECTL[#] is $\mathcal{B}^+(\mathcal{U}, \circ, \mathcal{U}\square)$.

We present a tree-style one-pass tableau for ECTL[#] continuing the analogous developments in linear-time case [11,12] and for CTL [11]. An indicative feature of this approach is a context-based tableau technique. Context-based tableaux have dual sequent calculi due to their handling of eventualities exclusively by using logical rules. To the best of our knowledge,

³ This notation goes back to [9], here we use its nice tuning by Nicolas Markey in [10]. In the last column we use a short CTL* formula $A\diamond(\circ p \wedge E\circ\neg p)$, not expressible by weaker logics. We found this formula indicative for CTL* as its validity is directly linked to the limit closure property [9].

BTL more expressive than CTL have not enjoyed the context-based tableau though other kinds of tableaux exist for these logics. There is a single-pass tableau for CTL that carries out an ‘on the fly’ eventualities checking (non-logical mechanism) following the Schwendimann’s approach [13]. For CTL*, which definitely is a super-logic of ECTL[#], different other kinds of tableau-style methods exist, remarkably [14–18]. Since CTL* is much more expressive than ECTL[#], such methods often utilise additional mechanisms (non only inference rules) to control loops, which are, for example, automata-theoretic-based mechanism [17]. This brings extra complexity, which is justified to handle the CTL* expressivity. However, simpler proofs could be obtained for a weaker logic such as ECTL[#]. There are also extensions of the tableau methods to super-logics of CTL*. For example, [19] introduces a two-pass tableau method for a logic that is a multiagent extension of CTL*. Tree-style one-pass tableaux (without additional procedures for checking meta-logical properties) have dual (cut-free) sequent calculi, see [12], enabling the construction of human-understandable proofs. In addition, these tableaux are well suited for the automation and are amenable for the implementation.⁴ Our tableau is effectively an AND-OR tree where nodes are labelled by sets of state (see the definitions in §2) formulae. There are difficult cases of ECTL[#] formulae that appear due to the enriched syntax: disjunctions of formulae in the scope of the A quantifier and conjunctions of formulae in the scope of the E quantifier. To tackle these cases, in addition to $\alpha - \beta$ rules, that are standard to the tableaux, we define novel β^+ -rules which use the context to force the eventualities to be fulfilled as soon as possible.

Outline of the paper. The rest of this paper, an extended version of [21], includes more examples, explanations, and detailed proofs of the results. It commences with §2 where we describe ECTL[#] as a sublogic of CTL*. The formulation of the tableau method is given in §3, where we define and explain tableau rules. A *systematic* tableau construction and relevant examples are introduced in §4. The soundness and completeness of our tableau method are proved in §5 and in §6, respectively; for the latter, we prove the refutational completeness and termination of the presented method. Finally, in §7 we draw the conclusions and prospects of future work that the presented results open.

2. The logic ECTL[#]

As ECTL[#] is a sublogic of CTL* we first recall CTL* syntax and semantics.

Definition 1 (*Syntax of CTL**). Given Prop is a fixed set of propositions, and $p \in \text{Prop}$, we define sets of state (σ) and path (π) CTL* formulae over Prop as follows: $\sigma ::= \top \mid p \mid \sigma_1 \wedge \sigma_2 \mid \neg\sigma \mid E\pi$ and $\pi ::= \sigma \mid \pi_1 \wedge \pi_2 \mid \neg\pi \mid \circ\pi \mid \pi \mathcal{U} \pi \mid \square\pi$. \square

In CTL*, and all BTL logics, well formed formulae are state formulae.

Definition 2 (*Labelled Kripke structure*). A Kripke structure, \mathcal{K} , is a triple (S, R, L) where $S \neq \emptyset$ is a set of states, $R \subseteq S \times S$ is a total binary relation, called *the transition relation*, and $L : S \rightarrow 2^{\text{Prop}}$ is a *labelling function*. \square

A *fullpath* x through a Kripke structure \mathcal{K} is an infinite sequence of states s_0, s_1, \dots such that $(s_i, s_{i+1}) \in R$, for every $i \geq 0$. Let ‘fullpaths(\mathcal{K})’ be the set of all fullpaths in \mathcal{K} . Given a fullpath $x = s_0, s_1, \dots, s_k, \dots$ ($k \geq 0$), we denote its state s_k by $x(k)$, its finite prefix by the sequence $x^{\leq k} = s_0, s_1, \dots, s_k$ and the suffix path $x^{\geq k} = s_k, s_{k+1}, \dots$. When a fullpath x is given, instead of $x(k)$ we will often write k , referring to k as ‘a state index of x ’. If x is a fullpath and y is a path such that $y(0) = x(k)$, for some $k > 0$, then the juxtaposition $x^{\leq k}y$ is a fullpath. Our Kripke structures are labelled directed graphs that correspond to Emerson’s R-generable structures, i.e. the transition relation R is suffix, fusion and limit closed [9]. For any \mathcal{K} , any $x \in \text{fullpaths}(\mathcal{K})$ and any natural number i , the notation $\mathcal{K} \upharpoonright x(i)$ denotes a Kripke structure with the set of states of \mathcal{K} restricted to those that are R -reachable from $x(i)$.

Definition 3. Given the structure $\mathcal{K} = (S, R, L)$, the relation \models , which evaluates path formulae in a given path x and state formulae at the state index i of the given path x , is defined below:

| | | |
|------------------------------------------------------|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\mathcal{K}, x, i \models \top$ | and | $\mathcal{K}, x, i \models p$ iff $p \in L(x(i))$. |
| $\mathcal{K}, x, i \models \neg\sigma$ | iff | $\mathcal{K}, x, i \models \sigma$ does not hold. |
| $\mathcal{K}, x, i \models \sigma_1 \wedge \sigma_2$ | iff | $\mathcal{K}, x, i \models \sigma_1$ and $\mathcal{K}, x, i \models \sigma_2$. |
| $\mathcal{K}, x, i \models E\pi$ | iff | there exists a path $y \in \text{fullpaths}(\mathcal{K} \upharpoonright x(i))$ such that $\mathcal{K}, y \models \pi$. |
| $\mathcal{K}, x \models \circ\pi$ | iff | $\mathcal{K}, x^{\geq 1} \models \pi$. |
| $\mathcal{K}, x \models \neg\pi$ | iff | $\mathcal{K}, x \models \pi$ does not hold. |
| $\mathcal{K}, x \models \pi_1 \wedge \pi_2$ | iff | $\mathcal{K}, x \models \pi_1$ and $\mathcal{K}, x \models \pi_2$. |
| $\mathcal{K}, x \models \pi_1 \mathcal{U} \pi_2$ | iff | there exists $k \geq i$ such that $\mathcal{K}, x^{\geq k} \models \pi_2$ and $\mathcal{K}, x^{\geq j} \models \pi_1$ for all $j \in \{0, \dots, k-1\}$. |
| $\mathcal{K}, x \models \square\pi$ | iff | $\mathcal{K}, x^{\geq j} \models \pi$ for all $j \geq 0$. |

In addition, for any set Σ of state formulae, $\mathcal{K}, x, i \models \Sigma$ iff $\mathcal{K}, x, i \models \sigma$, for all $\sigma \in \Sigma$. \square

⁴ An excellent survey of the seminal tableau techniques for temporal logics can be found in [20].

Many other usual operators can be derived from those introduced, in particular, the ‘falsehood’ constant $\mathbf{F} \equiv \neg\top$, and the disjunction operator $\varphi_1 \vee \varphi_2 \equiv \neg(\neg\varphi_1 \wedge \neg\varphi_2)$, as well as the temporal operator $\diamond\pi \equiv \top\mathcal{U}\pi$ and the universal path quantifier $A\pi \equiv \neg E\neg\pi$. It is also known that $\Box\pi \equiv \neg\diamond\neg\pi$ but, for technical convenience, we define it as a primitive operator. Let us recall some meta-logical concepts that are essential for the paper.

Definition 4 (*Syntactically consistent set of formulae*). A set Σ of state formulae σ is *syntactically consistent* abbreviated as Σ_{\top} if $\mathbf{F} \notin \Sigma$ and $\{\sigma, \neg\sigma\} \not\subseteq \Sigma$ for any σ ; otherwise, Σ is *inconsistent* denoted as Σ_{\perp} . \square

Definition 5 (*Satisfiability*). For a set of state formulae Σ , the set of its models, $\text{Mod}(\Sigma)$, is formed by all triples (\mathcal{K}, x, i) such that $\mathcal{K}, x, i \models \Sigma$. Σ is *satisfiable* ($\text{Sat}(\Sigma)$) if $\text{Mod}(\Sigma) \neq \emptyset$, otherwise Σ is *unsatisfiable* ($\text{UnSat}(\Sigma)$). \square

If $\text{Mod}(\Sigma) = \text{Mod}(\Sigma')$ then Σ and Σ' are equivalent denoted as $\Sigma \equiv \Sigma'$. For a set of state formulae Σ , if for any fullpath $x \in \text{fullpaths}(\mathcal{K})$, we have $\mathcal{K}, x, 0 \models \Sigma$, then we simply write $\mathcal{K} \models \Sigma$.

Definition 6 (*Cyclic sequence, cyclic path and cyclic Kripke structure*). Let z be a finite sequence of states $z = s_0, s_1, \dots, s_j$ such that, for every $0 \leq k < j$, $(s_k, s_{k+1}) \in R$. Then, z is *cyclic* iff there exists s_i , $0 \leq i \leq j$ such that $(s_j, s_i) \in R$. Let z be a finite cyclic sequence, the subsequence s_i, \dots, s_j of z is called a *loop* and s_i is called the *cyclining element*. We denote the loop as $\langle s_i, \dots, s_j \rangle^{\omega}$. A *cyclic path* over z is an infinite sequence $\text{path}(z) = s_0, s_1, \dots, s_{i-1} \langle s_i, s_{i+1}, \dots, s_j \rangle^{\omega}$.⁵ A Kripke structure \mathcal{K} is *cyclic* if every fullpath is a cyclic path over a cyclic sequence of states. \square

The fact that CTL* satisfiability can be reduced to the emptiness problem for automata on infinite trees (see [22,23]), ensures that the (non-empty) collection of models of a given satisfiable CTL* formula can be obtained by infinitely unwinding (in any possible way) a finite graph. Hence, for any CTL* formula ϕ , such that $\text{Mod}(\phi) \neq \emptyset$, there always exists a model $\mathcal{K} \in \text{Mod}(\phi)$ such that \mathcal{K} is cyclic. Therefore, when speaking about the satisfiability in CTL* (hence ECTL#) we can consider cyclic Kripke structures.

Proposing a new logic, ECTL#, we aim at defining a sublogic of CTL* that extends the ECTL+ formulae $\Box\diamond\sigma$ and $\diamond\Box\sigma$ (where σ means state formula), respectively, to $\Box(\sigma\mathcal{U}\sigma)$ and $\sigma\mathcal{U}\Box\sigma$.

Definition 7 (*Syntax of ECTL#*). The set of ECTL# formulae, over a fixed set of propositions Prop , are formed according to the following restriction of the CTL* grammar in Definition 1 for path formulae (state formulae are the same): $\pi ::= \sigma \mid \pi_1 \wedge \pi_2 \mid \neg\pi \mid \circ\sigma \mid \sigma\mathcal{U}\sigma \mid \Box\sigma \mid \sigma\mathcal{U}(\Box\sigma) \mid \Box(\sigma\mathcal{U}\sigma)$. \square

Note that the nesting of pure path formulae, totally unrestricted in CTL*, is now restricted by the grammar cases: $\circ\sigma \mid \sigma\mathcal{U}\sigma \mid \Box\sigma \mid \sigma\mathcal{U}\Box\sigma \mid \Box(\sigma\mathcal{U}\sigma)$. In particular, $a\mathcal{U}\Box(b \wedge \Box c)$ (where $a, b, c \in \text{Prop}$) is not an ECTL# formula because $b \wedge \Box c$ is directly in the scope of the \Box but is not a state formula. For technical convenience, we assume that the tableau construction applies to the formulae in *negation normal form* (shortly, nnf). Therefore, we introduce here a grammar for the set of ECTL# formulae that is closed under negation and requires the negation to apply to atomic propositions (instead of state and path formulae).

Definition 8 (*Syntax of ECTL# in nnf*). Let Prop be a fixed set of propositions, let $\rho \in \text{Prop}$ and let $\text{Lit} ::= \mathbf{F} \mid \top \mid \rho \mid \neg\rho$, be the set of literals. The set $\mathcal{F}_{\text{Prop}}$ of ECTL# formulae in nnf (over Prop) is given by the grammar:

$$\begin{aligned} \sigma &::= \text{Lit} \mid \sigma_1 \wedge \sigma_2 \mid \sigma_1 \vee \sigma_2 \mid E\pi \mid A\pi \\ \pi &::= \pi_1 \wedge \pi_2 \mid \pi_1 \vee \pi_2 \mid \circ\sigma \mid \sigma\mathcal{U}(\Box\sigma) \mid \Box(\sigma\mathcal{U}\sigma) \mid \Box(\sigma \vee \Box\sigma) \mid \sigma\mathcal{U}(\sigma \wedge \diamond\sigma) \end{aligned}$$

where σ means a state formula, π means a path formula, and $\diamond\sigma$ abbreviates $\top\mathcal{U}\sigma$. \square

The modified grammar is obtained by extending the state formulae grammar by $A\pi$ -formulae and the path formulae grammar by $\Box(\sigma \vee \Box\sigma)$ and $\sigma\mathcal{U}(\sigma \wedge \diamond\sigma)$. Cases $\sigma\mathcal{U}\sigma$ and $\Box\sigma$ are omitted because they respectively abbreviate $\sigma\mathcal{U}(\sigma \wedge \diamond\top)$ and $\Box(\sigma \vee \mathbf{F})$. Note that, for $a, b, c \in \text{Prop}$, the formula $\Box(a \vee \Box(b \vee \Box c))$ is not in $\mathcal{F}_{\text{Prop}}$ because $b \vee \Box c$ is not a state formula. The following proposition ensures that the set $\mathcal{F}_{\text{Prop}}$ is closed under negation.

Proposition 9 (*Closure under negation*). For any $\varphi \in \mathcal{F}_{\text{Prop}}$, we also have $\text{nnf}(\neg\varphi) \in \mathcal{F}_{\text{Prop}}$. Moreover, the negation of a state (resp. path) formula is a state (resp. path) formula.

⁵ Cyclic paths are also known as *ultimately periodic* paths.

Proof. By structural induction on the formulae, using the following equivalences (and well known classical ones):

1. $\neg A\varphi \equiv E\neg\varphi$
2. $\neg E\varphi \equiv A\neg\varphi$
3. $\neg\circ\varphi \equiv \circ\neg\varphi$
4. $\neg\Box\varphi \equiv \Diamond\neg\varphi$
5. $\neg\Box(\varphi_1 \mathcal{U} \varphi_2) \equiv \Box\neg\varphi_2 \vee \Diamond((\neg\varphi_1) \wedge (\neg\varphi_2))$
6. $\neg(\varphi_1 \mathcal{U} \Box\varphi_2) \equiv (\Box\Diamond\neg\varphi_2) \vee \Diamond(\neg\varphi_1 \wedge \Diamond\neg\varphi_2)$
7. $\neg(\varphi_1 \mathcal{U} (\varphi_2 \wedge \Diamond\varphi_3)) \equiv \Box(\neg\varphi_2 \vee \Box\neg\varphi_3) \vee ((\neg\varphi_2) \mathcal{U} (\neg\varphi_1 \wedge \neg\varphi_2))$

Equivalences 1-5 are very well known (e.g. [9]); the validity of 6 and 7 is easily established. It is also easy to see that 7, when φ_3 is τ , is reduced to the known equivalence $\neg(\varphi_1 \mathcal{U} \varphi_2) \equiv (\Box\neg\varphi_2) \vee (\neg\varphi_2 \mathcal{U} (\neg\varphi_1 \wedge \neg\varphi_2))$. \square

For simplicity, we will write $\neg\varphi$ instead of $\text{nnf}(\neg\varphi)$. Thus, $\neg A(p \mathcal{U} \Box q)$ represents $(E\Box\Diamond\neg q) \vee E\Diamond(\neg p \wedge \Diamond\neg q)$. Also, for a finite set $\Delta = \{\varphi_1, \dots, \varphi_n\}$, we let $\text{nnf}(\neg \bigwedge_{i=1}^n \varphi_i) = \neg\Delta$.

| Type of a difficult case | A-disjunctive formula | E-conjunctive formula |
|--------------------------|--------------------------|-----------------------------------------------|
| Example | $A(\circ q \vee \Box r)$ | $E(\circ r \wedge q \mathcal{U} \Box \neg p)$ |
| Our representation | $A(\circ q, \Box r)$ | $E(\circ r, q \mathcal{U} \Box \neg p)$ |

Fig. 2. Difficult cases of temporal operators in the scope of path quantifiers.

For ECTL[#], we identify the following difficult cases of the nesting and Boolean combinations of temporal operators in the scope of path quantifiers: A-disjunctive formula – disjunctions of temporal operators in the scope of A and E-conjunctive formula – conjunctions of temporal operators in the scope of E. For convenience, we will, respectively, write $A(\pi_1, \dots, \pi_n)$ and $E(\pi_1, \dots, \pi_n)$, where $n \geq 1$, and if “ \vee ” is in the scope of A it means \vee while being in the scope of E it means \wedge . Formulae serving as relevant examples in Fig. 2 will be used to illustrate tableau, in Fig. 6. Note that any A-formula (E-formula) σ can be transformed into an equivalent Boolean combination of A-disjunctive formulae $A(\pi_1, \dots, \pi_n)$ (E-conjunctive formulae $E(\pi_1, \dots, \pi_n)$), such that every π_i ($1 \leq i \leq n$) is of one of the following: $\circ\sigma$, $\sigma \mathcal{U} (\sigma \wedge \Diamond\sigma)$, $\sigma \mathcal{U} \Box\sigma$, $\Box(\sigma \vee \Box\sigma)$, and $\Box(\sigma \mathcal{U} \sigma)$, and σ stands for a state formula. For example, $A(((\circ q) \wedge (\Box E \circ r)) \vee \circ p)$ is equivalent to $A(\circ q) \wedge A(\Box E \circ r, \circ p)$; and $E(((\circ A \circ r) \vee (q \mathcal{U} \Box E \neg p)) \wedge \circ q)$ is equivalent to $E(\circ A \circ r) \vee E(q \mathcal{U} \Box E \neg p, \circ q)$. In what follows, Q abbreviates either of the path quantifiers. For a set of path formulae $\Pi = \{\pi_1, \dots, \pi_n\}$, we write $Q\Pi$ to denote $Q(\pi_1, \dots, \pi_n)$, and $Q\circ\Pi$ to denote $Q(\circ\pi_1, \dots, \circ\pi_n)$. If Φ is an empty set of formulae it means τ when Φ occurs in a conjunctive expression, and \mathbf{F} in a disjunctive expression. In particular, when Π is \emptyset then $A\Pi$ is \mathbf{F} and $E\Pi$ is τ . We write Σ, σ to represent the set $\Sigma \cup \{\sigma\}$. We consider that every formula $\sigma \in \mathcal{F}_{\text{Prop}}$ is given in its equivalent ‘negation normal form’, $\text{nnf}(\sigma)$.

3. The tableau method

3.1. Preliminaries

Definition 10 (Tableau, consistent node, closed branch). A tableau for a set of ECTL[#] state formulae Σ is a labelled tree T , where nodes are τ -labelled with sets of state formulae, such that the following two conditions hold:

- (a) The root is labelled by the set Σ .
- (b) Any other node m is labelled with sets of state formulae as the result of the application of one of the rules in Figs. 3, 4, 5 and 7 to its parent node n . Given the applied rule is R , we term m an R -successor of n .

A node n of T is consistent, abbreviated as n_{\top} , if $\tau(n)$ is a syntactically consistent set of formulae (see Def. 4), else n is inconsistent, abbreviated as n_{\perp} . If a branch b of T , contains $n_{\perp} \in b$, then b is closed else b is open. \square

To make the presentation more transparent we give an informal overview of the tableau construction. Any tableau has a root-node that is exclusively labelled by a set of state formulae. To extend a node we apply one of α , β or β^+ rule. The first two types of rules are standard to the tableau, and are essentially based on the fixpoint characterisation of $Q\Box$ and QU modalities, while β^+ rules are characteristic (and crucial!) for our construction. They tackle difficult cases of formulae in ECTL[#], and are related to our dedicated account of the eventualities. Namely, we treat an eventuality as occurring in some context, which, in turn, is a collection of all state formulae, called ‘an outer context’ or path formulae called ‘an inner context’. As we will see, β^+ rules use the context to force eventualities to be fulfilled as soon as possible. The $\alpha - \beta - \beta^+$ rules apply repeatedly until they produce an inconsistent node n_{\perp} , or a node with the labels that already occurred within the path under consideration. In the former case the expansion of the given branch terminates with n_{\perp} as its leaf. In the latter case, a repetitive node in the branch suggests that the input formula is satisfied forever, and we select another eventuality (if any) see §4.1. Obviously, n_{\perp} has an unsatisfiable $\tau(n)$ and is a ‘deadlock’ in the construction of a model. However, open branches do not necessarily give us a model. In particular, an open branch could be a prefix of a closed one. Later we introduce the notion of an expanded branch that enables the model construction. Once no more expansion rules

| | α | S_α |
|------------------------|---------------------------------------------|-------------------------------------------------------------------------------------------|
| (\wedge) | $\sigma_1 \wedge \sigma_2$ | $\{\sigma_1, \sigma_2\}$ |
| ($E\sigma$) | $E(\sigma_1, \dots, \sigma_n, \Pi)$ | $\{\sigma_1, \dots, \sigma_n, E\Pi\}$ |
| ($E\Box\mathcal{U}$) | $E(\Box(\sigma_1\mathcal{U}\sigma_2), \Pi)$ | $\{E(\sigma_1\mathcal{U}\sigma_2, \circ\Box(\sigma_1\mathcal{U}\sigma_2), \Pi)\}$ |
| ($A\Box\mathcal{U}$) | $A(\Box(\sigma_1\mathcal{U}\sigma_2), \Pi)$ | $\{A(\sigma_1\mathcal{U}\sigma_2, \Pi), A(\circ\Box(\sigma_1\mathcal{U}\sigma_2), \Pi)\}$ |

Fig. 3. ALPHA RULES. (Notation: σ, σ_i stand for state formulae and Π is a set of path formulae, possibly empty.)

| β_Rule | β | k | $S_{\beta_i} (1 \leq i \leq k)$ |
|--------------------------|-----------------------------------------------------------------|---------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (\vee) | $\sigma_1 \vee \sigma_2$ | 2 | $S_{\beta_1} = \{\sigma_1\}$ $S_{\beta_2} = \{\sigma_2\}$ |
| ($A\sigma$) | $A(\sigma_1, \dots, \sigma_n, \Pi)$ | $n + 1$ | $S_{\beta_1} = \{\sigma_1\}$ \vdots $S_{\beta_n} = \{\sigma_n\}$ $S_{\beta_{n+1}} = \{A\Pi\}$ |
| ($E\Box\sigma$) | $E(\Box(\sigma_1 \vee \Box\sigma_2), \Pi)$ | 2 | $S_{\beta_1} = \{\sigma_1, E(\circ\Box(\sigma_1 \vee \Box\sigma_2), \Pi)\}$ $S_{\beta_2} = \{-\sigma_1, \sigma_2, E(\circ\Box\sigma_2, \Pi)\}$ |
| ($E\mathcal{U}\sigma$) | $E(\sigma_1\mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3), \Pi)$ | 2 | $S_{\beta_1} = \{\sigma_2, E(\Diamond\sigma_3, \Pi)\}$ $S_{\beta_2} = \{\sigma_1, E(\circ(\sigma_1\mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3)), \Pi)\}$ |
| ($E\mathcal{U}\Box$) | $E(\sigma_1\mathcal{U}\Box\sigma_2, \Pi)$ | 2 | $S_{\beta_1} = \{E(\Box\sigma_2, \Pi)\}$ $S_{\beta_2} = \{\sigma_1, E(\circ(\sigma_1\mathcal{U}\Box\sigma_2), \Pi)\}$ |
| ($E\Box\mathcal{U}$) | $E(\Box(\sigma_1\mathcal{U}\sigma_2), \Pi)$ | 2 | $S_{\beta_1} = \{\sigma_2, E(\circ\Box(\sigma_1\mathcal{U}\sigma_2), \Pi)\}$ $S_{\beta_2} = \{\sigma_1, E(\circ\Box(\sigma_1\mathcal{U}\sigma_2), \Pi)\}$ |
| ($A\Box\sigma$) | $A(\Box(\sigma_1 \vee \Box\sigma_2), \Pi)$ | 3 | $S_{\beta_1} = \{\sigma_1, A(\circ\Box(\sigma_1 \vee \Box\sigma_2), \Pi)\}$ $S_{\beta_2} = \{-\sigma_1, \sigma_2, A(\circ\Box\sigma_2, \Pi)\}$ $S_{\beta_3} = \{A\Pi\}$ |
| ($A\mathcal{U}\sigma$) | $A(\sigma_1\mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3), \Pi)$ | 3 | $S_{\beta_1} = \{\sigma_2, A(\Diamond\sigma_3, \Pi)\}$ $S_{\beta_2} = \{\sigma_1, A(\circ(\sigma_1\mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3)), \Pi)\}$ $S_{\beta_3} = \{A\Pi\}$ |
| ($A\mathcal{U}\Box$) | $A(\sigma_1\mathcal{U}\Box\sigma_2, \Pi)$ | 2 | $S_{\beta_1} = \{A(\Box\sigma_2, \Pi)\}$ $S_{\beta_2} = \{\sigma_1, \sigma_2, A(\circ(\sigma_1\mathcal{U}\Box\sigma_2), \Pi)\}$ |

Fig. 4. BETA RULES. (Notation: σ, σ_i stand for state formulae, π_i stand for path formulae, and Π is a (possibly empty) set of path formulae.)

are applicable to the given branch with the last node n_\top , we are ensured that $\tau(n) = \Sigma, A\circ\Phi_1, \dots, A\circ\Phi_n, E\circ\Psi_1, \dots, E\circ\Psi_m$, where Σ is a set of literals. This labelling $\tau(n)$ is similar to a ‘state’ in the standard temporal tableau. Then the ‘next-state’ rule applies to generate the successors of n with the labels that are arguments of all $Q\circ$ modalities. The whole cycle of applying $\alpha - \beta - \beta^+$ and ‘next-state’ rules is repeated until the tableau construction terminates. The nature of our rules ensures that the terminated tableau represents a model for the tableau input if all the leaves in a collection of branches, called a bunch, are consistent and all eventualities occurring in looping branches are fulfilled, otherwise, the tableau input is unsatisfiable.

$$(Q\circ) \frac{\Sigma, A\circ\Phi_1, \dots, A\circ\Phi_\ell, E\circ\Psi_1, \dots, E\circ\Psi_k}{A\Phi_1, \dots, A\Phi_\ell, E\Psi_1 \& \dots \& A\Phi_1, \dots, A\Phi_\ell, E\Psi_k}$$

Fig. 5. NEXT-STATE RULE. (Notation: Σ is a (possibly empty) set of literals, and Φ_i, Ψ_i are non-empty sets of formulae.)

3.2. Alpha, beta rules and next-state rule

The α - and β -rules are the most elementary rules of our tableau system. An α -rule enlarges a branch with a node labelled by Σ, α , by a successor node labelled by Σ, S_α , where S_α is the set of formulae associated with α in Fig. 3. An α -rule is represented as $\frac{\Sigma, \alpha}{\Sigma, S_\alpha}$ while β -rules as $\frac{\Sigma, \beta}{\Sigma, S_{\beta_1} \mid \dots \mid \Sigma, S_{\beta_k}}$. β -rule splits a branch containing a node labelled by a set Σ, β (where β is one of the formulae of Fig. 4), in k new nodes each labelled by the corresponding Σ, S_{β_i} , according to Fig. 4. The next-state rule (Q \circ), Fig. 5, also splits the branch into k branches each of them rooted by node n labelled by a set $A\Phi_1, \dots, A\Phi_\ell, E\Psi_i$, for $i \in \{1, \dots, k\}$. This is the only rule of our system that splits branches in a conjunctive way. We use the symbol $\&$ to represent the generation of AND-successors of node n . When $\ell = k = 0$, the rule yields a unique new node labelled by the empty set. We assume that whenever $k = 0$ and $\ell > 0$, there exists a unique descendant labelled by $A\Phi_1, \dots, A\Phi_\ell$.

Example 11. Let n be a node such that $\tau(n) = \{a, \neg b, A\circ c, E\circ p, E\circ\neg p, A\circ\Box((E\circ p) \wedge (E\circ\neg p))\}$. Then the next-state rule (Q \circ) applies to n generating the following AND-successors of n : $\{Ac, p, A\Box((E\circ p) \wedge (E\circ\neg p))\}$ and $\{Ac, \neg p, A\Box((E\circ p) \wedge (E\circ\neg p))\}$. Note that Ac requires the application of the β -rule ($A\sigma$) to be reduced to c . \square

3.3. The uniform tableau

In this subsection we explain how to construct a tableau where leaves are labelled by sets of formulae of a specific form – *Uniform sets of state formulae*.

Definition 12 (*Elementary set of ECTL[#] state formulae*). A set of ECTL[#] state formulae is *elementary* if and only if it is exclusively formed by literals and formulae of the form $Q\circ\Pi$. \square

Proposition 13. Any set of ECTL[#] state formulae has a tableau T such that all its leaves are labelled by elementary sets of state formulae.

Proof. Repeatedly apply to every expandable node any applicable α -rule or β -rule. \square

Example 14. Fig. 6 depicts a tableau with elementary leaves for $A(\circ q, \Box r)$, $E(\circ r, q\mathcal{U}\Box\neg p)$, $E\circ q$. Recall that $A(\circ q, \Box r)$ is an abbreviation of $A(\circ q, \Box(r \vee \Box F))$.

Definition 15 (*Basic path/state formula and uniform set of formulae*). Every ECTL[#] path formula of the type $\circ\sigma, \sigma_1\mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3), \sigma_1\mathcal{U}(\Box\sigma_2), \Box(\sigma_1 \vee \Box\sigma_2), \Box(\sigma_1\mathcal{U}\sigma_2)$ is called *basic*. Every state formula $Q\Pi$ where Π is a set of basic-path formulae is also called *basic*. A set of state formulae Σ is *uniform* iff Σ is exclusively formed by literals and basic state formulae, and Σ contains at most one E-conjunctive formula. \square

Proposition 16. Any set of ECTL[#] state formulae Σ has a tableau T such that labels of all its leaves are uniform sets of state formulae. Moreover, each open branch of T contains exactly one application of (Q \circ).

Proof. Use Proposition 13 to construct a tableau with all its leaves labelled by elementary sets of formulae. Then apply the rule (Q \circ), to any relevant node and, finally, repeatedly apply (to every expandable node) the rules (E σ), ($A\sigma$), (\wedge), and (\vee). \square

Definition 17 (*Uniform tableaux*). For any set Σ of ECTL[#] state formulae, the tableau for Σ provided by Proposition 16 is denoted $\text{Uniform_Tableau}(\Sigma)$. \square




Fig. 6. A tableau whose leaves are elementary.

Example 18. Constructing a uniform tableau for the set $\{A(\circ q, \square r), E(\circ r, q\mathcal{U}\square\neg p), E\circ q\}$, we first obtain the tableau in Fig. 6. Then we apply the (Q \circ) rule enlarging each of the four branches and producing the following eight leaves, left to right (we refer to the node by its labels):

1. $A(q, \square r), E(r, \square\neg p)$
2. $A(q, \square r), E q$
3. $A q, E(r, \square\neg p)$
4. $A q, E q$
5. $A(q, \square r), E(r, q\mathcal{U}\square\neg p)$
6. $A(q, \square r), E q$
7. $A q, E(r, q\mathcal{U}\square\neg p)$
8. $A q, E q$

Then we apply the rules (A σ) and (E σ): the first branch is split into $q, r, E\square\neg p$ and $A\square r, r, E\square\neg p$; the second into q and $A\square r, q$; the third yields only a child $q, r, E\square\neg p$; the fourth and the eighth yield only q ; the fifth is split into two nodes $q, r, E(q\mathcal{U}\square\neg p)$ and $A\square r, r, E(q\mathcal{U}\square\neg p)$; the sixth into q and $A\square r, q$; and the seventh yields the unique child $q, r, E(q\mathcal{U}\square\neg p)$. \square

| β^+ -Rule | Σ, β | k | $S_{\Sigma, \beta_i}^+ (1 \leq i \leq k)$ |
|---------------------------------|---------------------------------------------------------------------------|------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $(A\mathcal{U}\sigma)^+$ | $\Sigma, A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond\sigma_3), \Pi)$ | 3 | $S_{\Sigma, \beta_1}^+ = \{\sigma_2, A(\diamond\sigma_3, \Pi)\}$ $S_{\Sigma, \beta_2}^+ = \{\sigma_1, A(\circ((\sigma_1 \wedge (\neg\Sigma \vee \varphi_\Pi))\mathcal{U} (\sigma_2 \wedge \diamond\sigma_3)), \Pi)\}$ $S_{\Sigma, \beta_3}^+ = \{A\Pi\}$ |
| $(A\mathcal{U}\square)^+$ | $\Sigma, A(\sigma_1 \mathcal{U} \square\sigma_2, \Pi)$ | 2 | $S_{\Sigma, \beta_1}^+ = \{A(\square\sigma_2, \Pi)\}$ $S_{\Sigma, \beta_2}^+ = \{\sigma_1, A(\circ((\sigma_1 \wedge (\neg\Sigma \vee \varphi_\Pi \vee \sigma_2))\mathcal{U} \square\sigma_2), \Pi)\}$ |
| β^+ -Rule | Σ, β | k | $S_{\Sigma, \beta_i}^+ (1 \leq i \leq k)$ |
| $(E\mathcal{U}\sigma\square)^+$ | $\Sigma, E\Pi$ | $2n$ | S_{Σ, β_1}^+ \vdots $S_{\Sigma, \beta_i}^+ = \begin{cases} \{\sigma_2, E(\diamond\sigma_3, \Pi^{-i})\} & \text{if } \pi_i = \sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond\sigma_3) \\ \{E(\square\sigma_2, \Pi^{-i})\} & \text{if } \pi_i = \sigma_1 \mathcal{U} \square\sigma_2 \end{cases}$ \vdots S_{Σ, β_n}^+ $S_{\Sigma, \beta_{n+1}}^+$ \vdots $S_{\Sigma, \beta_{2i}}^+ = \begin{cases} \{\sigma_1, E(\circ((\sigma_1 \wedge \neg\Sigma)\mathcal{U} (\sigma_2 \wedge \diamond\sigma_3)), \Pi^{-i})\} & \text{if } \pi_i = \sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond\sigma_3) \\ \{\sigma_1, E(\circ((\sigma_1 \wedge \neg\Sigma)\mathcal{U} \square\sigma_2), \Pi^{-i})\} & \text{if } \pi_i = \sigma_1 \mathcal{U} \square\sigma_2 \end{cases}$ \vdots $S_{\Sigma, \beta_{2n}}^+$ |

Fig. 7. BETA-PLUS RULES. (Notation: σ, σ_i stand for state formulae, Σ is a (possibly empty) set of state formulae, Π is a (possibly empty) set of basic-path formulae. Formula φ_Π is defined in Definition 19. We denote by $\Pi_{\mathcal{U}}$ the set of all formulae in Π that have the forms $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond\sigma_3)$ and $\sigma_1 \mathcal{U} \square\sigma_2$. $\Pi_{\mathcal{U}}$ is enumerated as $\{\pi_1, \dots, \pi_n\}$ for $n \geq 1$, and $\Pi^{-i} = \Pi \setminus \{\pi_i\}$.)

3.4. The beta-plus rules

In this subsection we extend our set of tableau rules with the new four rules named as β^+ -rules (Fig. 7). These rules, similarly to β -rules, also split a branch, but this time into a number of branches depending on the treated formula. The rules for A-disjunctive formulae apply to a label Σ, β , where β has the form $A(\pi, \Pi)$, Π is a set of basic-path formulae, and π is either $\sigma \mathcal{U} (\sigma \wedge \diamond\sigma)$ or $\sigma \mathcal{U} \square\sigma$. The rule $(E\mathcal{U}\sigma\square)^+$ for E-conjunctive formulae applies to a set Σ, β , where β has the form $E\Pi$ and Π is a set of basic-path formulae that contains at least one formula $\sigma \mathcal{U} (\sigma \wedge \diamond\sigma)$ or $\sigma \mathcal{U} \square\sigma$. The β^+ -rules are the only rules in our system that make use of the so-called context for forcing the eventualities to be satisfied as soon as possible. The context is given by the sets Σ containing state formulae and Π containing path formulae. We name Σ the outer context and Π the inner context. The outer context is used by all the β^+ -rules. The inner context is only needed to deal with formulae $A\Pi$. The following formula, φ_Π , introduced in Definition 19 is used to manage the inner context Π in rules $(A\mathcal{U}\sigma)^+$ and $(A\mathcal{U}\square)^+$.

Definition 19 (Formula φ_Π for β^+ -rules on A-disjunctive formulae). Let Π be a set of basic path formulae. We define the formula φ_Π to be the following disjunction of state formulae (φ_Π is **F**, if the below disjunction is empty):

$$\bigvee_{\square(\sigma_1 \vee \square\sigma_2) \in \Pi} (\sigma_1 \vee \sigma_2) \vee \bigvee_{\sigma_1 \mathcal{U} \square\sigma_2 \in \Pi} \sigma_2 \vee \bigvee_{\square(\sigma_1 \mathcal{U} \sigma_2) \in \Pi} E(\diamond\sigma_2). \quad \square$$

It is worth noting that each β^+ -rule, when applied to some formula of the form $Q(\sigma_1 \mathcal{U} \varphi, \Pi)$ –where φ could be $\sigma_2 \wedge \diamond \sigma_3$ or $\square \sigma_2$ – generates one or more successors that contain a formula of the form $Q(\circ((\sigma_1 \wedge \sigma) \mathcal{U} \varphi), \Pi)$ where σ depends on both the inner and the outer context, and is defined depending on whether Q is E or A . We call $(\sigma_1 \wedge \sigma) \mathcal{U} \varphi$ the *next-step variant* of $\sigma \mathcal{U} \varphi$. Example 20 illustrates the main ideas behind the application of β^+ -rules $(A\mathcal{U}\sigma)^+$ and $(A\mathcal{U}\sigma)$.

Example 20. The β^+ -rules $(A\mathcal{U}\sigma)^+$ applies to one selected formula with exactly one marked eventuality. Consequently, the $(A\mathcal{U}\sigma)$ -rule applies to all the eventualities (in the selected formula) except to the marked one.




Fig. 8. Application of $(A\mathcal{U}\sigma)^+$ and $(A\mathcal{U}\sigma)$ rules to $\{-b, A(a\mathcal{U}b, p\mathcal{U}q)\}$.

In Fig. 8 the marked eventualities are in gray boxes. Assume $a\mathcal{U}b$ is the marked eventuality. Then, the $(A\mathcal{U}\sigma)^+$ -rule can be applied to $a\mathcal{U}b$ with outer context $\Sigma = \{-b\}$ and inner context, $\Pi = \{p\mathcal{U}q\}$. According to Definition 19, φ_Π is F . Therefore, $S_{\Sigma, \beta_1}^+ = \{b\}$, $S_{\Sigma, \beta_3}^+ = \{A(p\mathcal{U}q)\}$, and $S_{\Sigma, \beta_2}^+ = \{a, A(\circ((a \wedge b)\mathcal{U}b), p\mathcal{U}q)\}$ respectively generate nodes n_1 , n_2 and n_3 . In node n_2 , the $(A\mathcal{U}\sigma)$ -rule is applied to $p\mathcal{U}q$. That produces three new nodes. Regarding node n_3 , the marked eventuality disappears. Then, a new selection is made and $p\mathcal{U}q$ is marked. Consequently, $(A\mathcal{U}\sigma)^+$ -rule is applied with outer context $\Sigma = \{-b\}$. The inner context is the empty set. Note that all leaves in Fig. 8 are elementary. Hence, the construction of the tableau, would continue applying the next-state rule $(Q\circ)$. \square

3.5. Simplification rules

A large set of simplification rules can be used to reduce the tableau construction. Here we only mention those that are essential for termination. First, to stop the growth of the subformula σ in the successive next-step variants $(\sigma_1 \wedge \sigma) \mathcal{U} \varphi$, we use trivial simplification rules such as $\varphi \wedge \varphi \rightarrow \varphi$ and $\varphi \vee \varphi \rightarrow \varphi$, as well as classical subsumptions rules. Second, to simplify the detection of equal labels (for looping in tableau branches) we use the following (subsumption-based) rules:

$$\begin{aligned} (\square E \square \mathcal{U}) \quad & E(\sigma_1 \mathcal{U} \sigma_2, \square(\sigma_1 \mathcal{U} \sigma_2), \Pi) \rightarrow E(\square(\sigma_1 \mathcal{U} \sigma_2), \Pi). \\ (\square A \square \mathcal{U}) \quad & \text{If } \Pi' \subseteq \Pi \text{ then } A(\sigma_1 \mathcal{U} \sigma_2, \Pi) \wedge A(\square(\sigma_1 \mathcal{U} \sigma_2), \Pi') \rightarrow A(\square(\sigma_1 \mathcal{U} \sigma_2), \Pi'). \end{aligned}$$

Finally, to prevent the duplications of the original eventuality $\sigma_1 \mathcal{U} \sigma_2$ and its successive new-step variants by rules $(Q\square\mathcal{U})$ and $(Q\mathcal{U}\sigma)^+$, and to ensure termination, we use the following (subsumption-based) simplification rules:

$$\begin{aligned} (\square A \sigma \mathcal{U}) \quad & \sigma_2 \wedge A(\sigma_1 \mathcal{U} \sigma_2, \Pi) \rightarrow \sigma_2. \\ (\square E \mathcal{U} \sigma) \quad & E((\sigma_1 \wedge \sigma) \mathcal{U} \varphi, \sigma_1 \mathcal{U} \varphi, \Pi) \rightarrow E((\sigma_1 \wedge \sigma) \mathcal{U} \varphi, \Pi). \\ (\square A \mathcal{U} \sigma) \quad & \text{If } \Pi' \subseteq \Pi \text{ then } A((\sigma_1 \wedge \sigma) \mathcal{U} \varphi, \Pi') \wedge A(\sigma_1 \mathcal{U} \varphi, \Pi) \rightarrow A((\sigma_1 \wedge \sigma) \mathcal{U} \varphi, \Pi'). \end{aligned}$$

3.6. The role of φ_Π in the beta-plus rules

Let us consider a set of formulae $\Phi = \Sigma, A(\sigma \mathcal{U} \chi, \Pi)$. A model, \mathcal{K} , of Φ could satisfy $A(\sigma \mathcal{U} \chi, \Pi)$ because each fullpath of \mathcal{K} satisfies either $\sigma \mathcal{U} \chi$ or some formula $\pi \in \Pi$. The next-step variant of $\sigma \mathcal{U} \chi$ is $\circ(\sigma \wedge (\neg \Sigma \vee \varphi_\Pi)) \mathcal{U} \chi$, which makes $\neg \Sigma$ or φ_Π satisfiable before χ is satisfied. The former produces open branches where χ is fulfilled as soon as possible, whereas the latter produces open branches that satisfy some of the $\pi \in \Pi$. Therefore, φ_Π allows to generate a model from a branch in which $\sigma \mathcal{U} \varphi$ is not fulfilled, and some $\pi \in \Pi$ is satisfied. Example 21 illustrates these ideas from the constructive view, i.e. when we construct a tableau for a formula $A(\pi_1, \dots, \pi_n)$.

Example 21. Let $\Pi = \{\square c, r\mathcal{U}\square s, \square(p\mathcal{U}q)\}$, and consider an application of the rule $(A\mathcal{U}\sigma)^+$ to the formula $A(a\mathcal{U}b, \Pi)$, where $a, b, c, p, q, r, s \in \text{Prop}$ (see Fig. 9). The outer context, namely Σ , is empty and the inner context is Π . Then $\neg \Sigma$ is F and $\varphi_\Pi = c \vee s \vee E \diamond q$. Hence, the second child, namely S_{Σ, β_2}^+ , raised by the application of $(A\mathcal{U}\sigma)^+$ is labelled by $\{a, A(\circ \alpha_1, \Pi)\}$ where $\alpha_1 = (a \wedge \varphi_\Pi) \mathcal{U} b = (a \wedge (c \vee s \vee E \diamond q)) \mathcal{U} b$. Then, Uniform_Tableau applies the (corresponding) rules to $\square c, r\mathcal{U}\square s, \square(p\mathcal{U}q)$, and, finally, $(Q\circ)$ and the simplification rule $(\square A \square \mathcal{U})$, obtaining the node n_3 . Now, one of the leaves raised by Uniform_Tableau is the fifth node; by applying here $(Q\circ)$ and $(\square A \square \mathcal{U})$ we obtain the last node n_6 such that

Procedure *Eventuality_Selection* chooses formula $Q\Pi$ and if $Q = A$ then the procedure also marks one eventuality, according to the priorities of selection and marking in Definition 24. We denote by $\pi_{\mathcal{U}}$ the marked eventuality in the selected formula $A\Pi$.

Procedure *Apply_β⁺-rule(Σ)* applies the corresponding rules depending on the selected formula $Q\Pi$ and on the marked eventuality in the case $Q = A$:

- If $Q = A$ and $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3) \in \Pi$ is the marked eventuality, then apply $(A\mathcal{U}\sigma)^+$.
- If $Q = A$ and $\sigma_1\mathcal{U}\square\sigma_2 \in \Pi$ is the marked eventuality, then apply $(A\mathcal{U}\square)^+$.
- If $Q = A$ and $\square(\sigma_1\mathcal{U}\sigma_2) \in \Pi$ is the marked eventuality, then first apply the rule $(A\square\mathcal{U})$ and then the rule $(A\mathcal{U}\sigma)^+$ with the marked eventuality $\sigma_1\mathcal{U}\sigma_2$.
- If $Q = E$ and Π contains at least one $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ or one $\sigma_1\mathcal{U}\square\sigma_2$ then apply $(E\mathcal{U}\sigma\square)^+$.
- If $Q = E$ and Π contains at least one $\square(\sigma_1\mathcal{U}\sigma_2)$ (but none $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ and none $\sigma_1\mathcal{U}\square\sigma_2$), then first apply the rule $(E\square\mathcal{U})$ to every $\square(\sigma_1\mathcal{U}\sigma_2)$ and then the rule $(E\mathcal{U}\sigma\square)^+$.

Each application of a β^+ -rule on the selected $A\Pi$ introduces a next-step variant of the marked eventuality and each application of a β^+ -rule on the selected $E\Pi$ introduces a next-step variant for each $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ and each $\sigma_1\mathcal{U}\square\sigma_2$.

The call *Eventuality_Selection(Σ)* keeps the selection of the formula $E\Pi$ such that Π contains at least one $\sigma\mathcal{U}(\sigma \wedge \diamond\sigma)$ or $\sigma\mathcal{U}\square\sigma$, or keeps the selection of the formula $A\Pi \in \Sigma$ which contains the next-step variant of the marked eventuality, or selects a new formula $A\Pi \in \Sigma$ that contains an eventuality. The latter can happen for three reasons. When formulae $E\Pi$ do not contain any $\sigma\mathcal{U}(\sigma \wedge \diamond\sigma)$ nor $\sigma\mathcal{U}\square\sigma$, or there is no marked eventuality in formulae $A\Pi$, or when there is one, the node ℓ , ($\Sigma = \tau(\ell)$) is a loop-node (see Definition 25), and the branch from the root-node to ℓ is not *eventuality-covered* (see Definition 26). In this case, a new selection should be made because there are eventualities that have never been marked but they should be. This way we introduce the term *eventuality-covered*. When a branch is eventuality-covered, its leaf is a loop-node and we are sure that, along the loop, at least some eventuality in each A-disjunctive formula and all eventualities in each E-conjunctive formula have been fulfilled. Consequently, the branch is an expanded open branch (see Definition 37) and represents a path in a possible model. It is worth mentioning that the only requirement for a branch to be eventuality-covered is to mark all necessary eventualities. The fact that formulae $E\Pi$ are kept selected whereas they contain some eventuality and formulae $A\Pi$ are kept selected where the next-step variant of the marked eventuality is kept marked ensures that every eventuality in $E\Pi$ and at least one in each $A\Pi$ is fulfilled.

When making the selection, priorities are used as stated in Definition 24. The idea behind priorities is that a tableau branch represents a path in a cyclic Kripke structure that is a possible model for the input formula. Therefore, it consists of a (possibly empty) initial sequence of states followed by a looping-sequence. Selectable formulae are classified into two sets - those of the highest priority and those of the lowest priority. The non-looping initial sequence is the first part of the model, hence we firstly select formulae $A\Pi$ where Π is exclusively formed by formulae of the form $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ and formulae $E\Pi$ where Π contains at least one eventuality of the form $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ or $\sigma_1\mathcal{U}\square\sigma_2$. These are the highest priority formulae, which cannot produce a loop. When one of the former formulae $A\Pi$ is selected, one of the $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ is marked, namely π . By means of the rule $(A\mathcal{U}\sigma)^+$, in a finite number of steps, either the branch close or π is either fulfilled (note that in the first branch $A(\diamond\sigma_3, \Pi)$ is also of the highest priority) or deleted (the third branch of $(A\mathcal{U}\sigma)^+$). In any case the original formulae $A\Pi$ disappears. When one of the latter formulae $E\Pi$ is selected, the successive applications of the rule $(E\mathcal{U}\sigma\square)^+$ ensure (excluding the case when the branch closes) the fulfilment of all $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ or $\sigma_1\mathcal{U}\square\sigma_2$ in a finite number of states. Note that $E(\diamond\sigma_3, \Pi)$ is also of the highest priority. Once such formulae are fulfilled, all formulae $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ have disappeared from the E-conjunctive formula, whereas $\square\sigma_2$ remains in the conjunction for all $\sigma_1\mathcal{U}\square\sigma_2 \in \Pi$. Hence, the residual $E\Pi'$ is of the lowest-priority. On the contrary, the lowest priority formulae could produce a loop. They are formulae $A\Pi$ where Π contains at least one $\sigma_1\mathcal{U}\square\sigma_2$ or $\square(\sigma_1\mathcal{U}\sigma_2)$ and formulae $E\Pi$ where Π contains at least one $\square(\sigma_1\mathcal{U}\sigma_2)$ (but are not of the highest priority). They could produce a loop in a finite number of steps, since the subformulae starting by \square remains forever in the E-disjunctive formulae, whereas in the A-disjunctive formulae they can either remain or disappear. In the latter case, the residual A-disjunctive formulae could become non-selectable. It is easy to see that non-selectable formulae necessarily produce a loop.

Example 23. Consider $\Sigma_0 = \{A(a\mathcal{U}b, b\mathcal{U}c), E(p\mathcal{U}q, \square(r\mathcal{U}s)), A\square(c\mathcal{U}d), \neg b, A\square e\}$. The first two formulae have the highest-priority, the third has the lowest priority, and the last two are non-selectable. Suppose that we select $A(a\mathcal{U}b, b\mathcal{U}c)$ and mark $a\mathcal{U}b$, since $\neg b \in \Sigma_0$, the left-most open branch of rule $(A\mathcal{U}\sigma)^+$ contains $a, A(\circ((a \wedge \neg\Sigma'_0)\mathcal{U}b), b\mathcal{U}c)$ where $\Sigma'_0 = \Sigma_0 \setminus \{A(a\mathcal{U}b, b\mathcal{U}c)\}$. After applying the corresponding α and β rules to the remaining formulae, the first stage s_0 (the first state of the model) contains the atoms $\{a, q, s, d, e\}$. Then, by the next-step rule $(Q\circ)$, the first node of the second stage s_1 is $\Sigma_1 = \{A((a \wedge \neg\Sigma'_0)\mathcal{U}b, b\mathcal{U}c), E\square(r\mathcal{U}s), A\square(c\mathcal{U}d), A\square e\}$ where the first formula is kept selected and the first eventuality is kept marked. Note that the second formula has now the lowest priority. Then we apply $(A\mathcal{U}\sigma)^+$ to the first formulae and the corresponding α and β rules to the remaining formulae in Σ_1 , generating the set of atoms in the left-most branch are $\{b, s, d, e\}$. Then, by the next-step rule $(Q\circ)$, the first node of the third stage s_2 is $\Sigma_2 = \{E\square(r\mathcal{U}s), A\square(c\mathcal{U}d), A\square e\}$ where the first two formulas are of the lowest priority and the third is non-selectable. By selecting the first formula, the atoms in the stages s_2 (of the left-most branch) are $\{s, d, e\}$ and the new uniform set at the first node of the following stage is $\Sigma_3 = \Sigma_2$. However, $A\square(c\mathcal{U}d)$ has not been selected inside the loop, hence we produce one stage more, s_3 , with atoms

$\{s, d, e\}$. Then, by $(Q\circ)$, we obtain $\Sigma_4 = \Sigma_2$ and the branch is eventuality-covered. Therefore, we have a model for Σ_0 is $s_0, s_1 \{s_2, s_3\}^\omega$. \square




Fig. 10. A branch in the systematic tableau for $A(p\mathcal{U}\neg p, \Box p), A\Box(a\mathcal{U}E\Box c)$.

Definition 24 (Priorities for eventuality selection). The selectable formulae of the highest priority for Eventuality_Selection are the formulae of the following two forms:

- $A\Pi$ where Π is exclusively formed by formulae of the form $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$.
- $E\Pi$ where Π contains at least one eventuality of the form $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ or $\sigma_1\mathcal{U}\Box\sigma_2$.

Consequently, the (selectable) formulae of the lowest priority are the formulae of the following two forms:

- $A\Pi$ where Π contains at least one $\sigma_1\mathcal{U}\Box\sigma_2$ or $\Box(\sigma_1\mathcal{U}\sigma_2)$.
- $E\Pi$ where Π does not contain any $\sigma_1\mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ nor $\sigma_1\mathcal{U}\Box\sigma_2$, and Π contains at least one $\Box(\sigma_1\mathcal{U}\sigma_2)$. \square

Once all the highest priority formulae have been selected in a branch, the only selectable formulae are the lowest priority formulae. At this point, the objective is to get a loop-node that makes the branch eventuality-covered.

Definition 25 (Loop-node). Let b be a tableau branch and $n_i \in b$ ($0 \leq i$). Then n_i is a loop-node if there exists $n_j \in b$ ($0 \leq j < i$) and $\tau(n_i) = \tau(n_j)$. We say that n_j is a companion node of n_i . \square

Definition 26 (Eventuality-covered branch). A tableau branch $b = n_0, n_1, \dots, n_i$ is eventuality-covered if n_i is a loop-node, with a companion node n_j ($0 \leq j < i$), both labelled by a uniform set Σ of non-selectable and the lowest priority formulae such that every selectable formula $Q\Pi \in \tau(n_i)$ is selected in some node n_k ($j \leq k < i$) and for every selected $A\Pi$ exactly one eventuality in Π is marked in some node n_k such that $j \leq k < i$. \square

The procedure `Eventuality_Selection` performs in some fair way that ensures that any open branch will ever be *eventuality-covered*.

Definition 27 (*Non-expandable node*). A node n is non-expandable, if $\tau(n) = \Sigma_{\perp}$ or n is a loop-node of branch b which is eventuality-covered. Otherwise, n is expandable. \square

Consequently, an expandable node is either a node that is not a loop-node or a loop-node whose branch is not eventuality-covered. It is worth noting that a formula of the lowest priority could be selected more than once in a branch because the loop-node could change along the branch. In the following Example 28, we illustrate this issue.

Example 28. Fig. 10 shows a branch in the systematic tableau for $\Sigma_0 = \{A(p\mathcal{U}\neg p, \Box p), A\Box(a\mathcal{U}E\Box c)\}$ where, for readability, the marked eventualities are in gray boxes. The call `Eventuality_Selection`(Σ_0) selects the formula $A(p\mathcal{U}\neg p, \Box p)$ in n_1 . Generating n_2 , when we apply the $(A\mathcal{U}\sigma)^+$ rule to $A(p\mathcal{U}\neg p, \Box p)$, the inner context is p and the outer context is $A\Box(a\mathcal{U}E\Box c)$. Hence, in the S_{Σ, β_2}^+ branch, the next-step variant of $p\mathcal{U}\neg p$ is $p \wedge (\neg(A\Box(a\mathcal{U}E\Box c) \vee p))\mathcal{U}\neg p$. By classical subsumption (included in our simplification rules), $p \wedge (\neg(A\Box(a\mathcal{U}E\Box c) \vee p)) \rightarrow p$, hence the next-step variant is $p\mathcal{U}\neg p$, and the formula $A(\circ(p\mathcal{U}\neg p), \Box p)$ is added to the current stage. Then, applying $(A\circ\sigma)$ to $A(\circ(p\mathcal{U}\neg p), \Box p)$, $(A\Box\mathcal{U})$ to $A\Box(a\mathcal{U}E\Box c)$, and $(A\mathcal{U}\sigma)$ to $A(a\mathcal{U}E\Box c)$, the node n_2 is obtained. After the application of $(Q\circ)$ and $(\Box A\Box\mathcal{U})$, n_3 is obtained. Node n_3 is a loop-node whose companion node is n_1 ($\tau(n_3) = \tau(n_1)$). However, the branch is not eventually-covered since the eventuality $a\mathcal{U}E\Box c$ is not selected inside the loop. Therefore, we obtain n_4 by the call `Eventuality_Selection`($(\tau(n_3))$) which selects $A(a\mathcal{U}E\Box c)$. After applying the $(A\mathcal{U}\sigma)^+$ rule to $A(a\mathcal{U}E\Box c)$ (once the $(A\Box\mathcal{U})$ rule is applied), the branch S_{Σ, β_1}^+ expands to n_5 . After that, `Uniform_Tableau` gets one expandable node labelled by the uniform set $\{E\Box c, A(p\mathcal{U}\neg p, \Box p), A\Box(a\mathcal{U}E\Box c)\}$ as represented in n_6 . The call `Eventuality_Selection`($(\tau(n_6))$) selects again the formula $A(p\mathcal{U}\neg p, \Box p)$. Now, the inner context is p and the outer context is $\{E\Box c, A\Box(a\mathcal{U}E\Box c)\}$. Hence, by subsumption, $p \wedge (\neg(E\Box c) \vee \neg(A\Box(a\mathcal{U}E\Box c)) \vee p) \rightarrow p$. Hence, the S_{Σ, β_2}^+ branch contains again the next-step variant $p\mathcal{U}\neg p$ in n_7 . Then, expanding the `Uniform_Tableau` we obtain n_8 which is an expandable loop-node because $\tau(n_8) = \tau(n_6)$. However, the branch is not yet eventually-covered since $a\mathcal{U}E\Box c$ has not been marked inside the loop. Then, the selected formula in n_9 is $A(a\mathcal{U}E\Box c)$. Finally, `Uniform_Tableau` obtains a non-expandable loop-node, thus the given branch is eventually-covered - depicted in Fig. 10 it represents $\{p, a\}\{p, c\}^w$, which is a model of Σ_0 . However, considering all the nodes in the branch, one would realize that the model represented is $\{p, a\}\{p, c\}\{p, a, c\}\{p, c\}^w$. \square

Definition 29 (*Bunch in a tableau, closed bunch and tableau*). A bunch b is a collection of branches that is maximal with respect to $(Q\circ)$ -successor, i.e. every $(Q\circ)$ -successor of any node in b is also in b . A bunch is closed iff at least one of its branches is closed, otherwise it is open. A tableau is closed iff all its bunches are closed. \square

Therefore, any open tableau has at least one open bunch, formed by one or more open branches. To complete, this section we provide two examples: a closed tableau and an open tableau. We mark eventualities in gray boxes and use large circles to represent the generation of AND-nodes or bunches.




Fig. 11. A closed tableau for $A(\tau\mathcal{U} p), E\Box\neg p$.

Example 30. Fig. 11 shows a closed tableau for $A(\tau\mathcal{U} p), E\Box\neg p$. Note that, in the two applications of the rule $(A\mathcal{U}\sigma)^+$, the inner context is empty and the outer context is $E\Box\neg p$ whose negation in nnf is $A\Diamond p$. Hence, the label of the rightmost node, $A(\circ((A\Diamond p) \wedge (A\Diamond p))\mathcal{U} p)$, is the simplification of the selected formula $A(\circ((A\Diamond p) \wedge (A\Diamond p))\mathcal{U} p)$. \square




Fig. 12. Open bunch in the tableau for $p, A\Box(E\Box p \wedge E\Box \neg p), A(\Diamond \neg p, \Box p)$ and represented model.

Example 31. On the left of Fig. 12 we depict a representative open bunch of a tableau for the set of formulae: $p, A\Box(E\Box p \wedge E\Box \neg p), A(\Diamond \neg p, \Box p)$. For saving space, we apply at once the Uniform_Tableau procedure subsequently choosing one of the leaves produced. Note that, for each node, we draw only one of the OR-children, but all the AND-children. In the marked eventuality, $\neg p \vee E\Diamond(A\Box \neg p \vee A\Box p) \vee p$ comes from the negation of the outer context, and the disjunct p from the inner context. By ‘Simplification’ $\neg p \vee E\Diamond(A\Box \neg p \vee A\Box p) \vee p$ is reduced to τ (in the left-hand child). In the label of the right-hand node, $\neg p$ subsumes $A(\dots)\mathcal{U}\neg p, \Box p$. This open bunch represents a model (of the input set of formulae) that we depict on the right of Fig. 12. \square

5. Soundness

To prove the soundness of our tableau method (Theorem 34) we show that every tableau rule preserves satisfiability (Lemma 33). To prove the latter we essentially use the limit closure property, ensuring that the satisfiability of the negated inner context is preserved from segments of a limit path to the limit path itself (Proposition 32). The use of φ_Π (Definition 19) is crucial here.

Proposition 32 (Preservation of the negated inner context). *Let Π be any set of basic path formulae and let φ_Π be as in Definition 19. Let \mathcal{K} be a Kripke structure, $x_1 \in \text{fullpaths}(\mathcal{K})$ such that $\mathcal{K}, x_1, 0 \models \neg\Pi$. Let $y = x_1^{\leq i_1} x_2^{\leq i_2} \dots x_k^{\leq i_k} \dots$ be a limit path in $\text{fullpaths}(\mathcal{K})$, for some $i_1 > 0$ and some $x_2^{\leq i_2} \dots x_k^{\leq i_k} \dots$. Then $\mathcal{K}, y, 0 \models \neg\pi$ holds for all $\pi \in \Pi$, provided that the following two conditions hold for all $n \geq 1$:*

- (a) $\mathcal{K}, x_1^{\leq i_1} x_2^{\leq i_2} \dots x_n, j \models \neg\sigma_2$ for all $\sigma_1 \mathcal{U}(\sigma_2 \wedge \Diamond \sigma_3) \in \Pi$ and all $j \in \{0..i_n\}$, and
- (b) $\mathcal{K}, x_1^{\leq i_1} x_2^{\leq i_2} \dots x_n^{\leq i_n}, i_n \models \neg\varphi_\Pi$.

Proof. We check the four cases of a basic path formula $\pi \in \Pi$. If π is of the form $\Box\sigma$, then $\mathcal{K}, y, 0 \models \neg\Box\sigma$ because $\mathcal{K}, y, 0 \models \neg\pi$ and $i_1 > 0$. If π is of the form $\sigma_1 \mathcal{U}(\sigma_2 \wedge \Diamond \sigma_3)$, then property (a) ensures that every state in y satisfies $\neg\sigma_2$. Therefore, $\neg(\sigma_1 \mathcal{U}(\sigma_2 \wedge \Diamond \sigma_3))$ is satisfied along the limit path y . For the remaining three cases, on the basis of (b) and Definition 19, we can prove the following three facts: (1) If $\pi = \Box(\sigma_1 \vee \Box\sigma_2)$, then $\mathcal{K}, x_1^{\leq i_1} x_2^{\leq i_2} \dots x_n^{\leq i_n}, i_n \models \neg\sigma_1 \wedge \neg\sigma_2$ for all n . (2) If $\pi = \Box(\sigma_1 \mathcal{U}\sigma_2)$, then we have that $\mathcal{K}, x_1^{\leq i_1} x_2^{\leq i_2} \dots x_n^{\leq i_n}, i_n \models \neg E(\Diamond \sigma_2)$ for all n . (3) If $\pi = \sigma_1 \mathcal{U}\Box\sigma_2$, then $\mathcal{K}, x_1^{\leq i_1} x_2^{\leq i_2} \dots x_n^{\leq i_n}, i_n \models \neg\sigma_2$ for all n . Therefore, in any of the three cases, we can ensure that $\mathcal{K}, y, 0 \models \pi$. \square

Lemma 33 (Soundness of the tableau rules). *For any set of state formulae Σ :*

- (i) For any α -formula α : $\text{Sat}(\Sigma, \alpha)$ iff $\text{Sat}(\Sigma, S_\alpha)$.
- (ii) For any β -formula β of range k : $\text{Sat}(\Sigma, \beta)$ iff $\text{Sat}(\Sigma, S_{\beta_i})$ for some $1 \leq i \leq k$.
- (iii) For any β^+ -formula β of range k : $\text{Sat}(\Sigma, \beta)$ iff $\text{Sat}(\Sigma, S_{\Sigma, \beta_i}^+)$ for some $1 \leq i \leq k$.
- (iv) If Σ is a set of consistent literals: $\text{Sat}(\Sigma, A\Box\Phi_1, \dots, A\Box\Phi_n, E\Box\Psi_1, \dots, E\Box\Psi_m)$ iff for all $0 \leq i \leq m$: $\text{Sat}(A\Phi_1, \dots, A\Phi_n, E\Psi_i)$. \square

Proof. Noting that (i), (ii) and (iv) can be easily proved by the ‘systematic’ application of the semantic definitions of temporal operators, we prove (iii). The ‘only if’ direction for each of the cases of β^+ -rules is trivial. We will prove the ‘if’ direction

of the three rules $(EU\sigma\Box)^+$, $(AU\sigma)^+$ and $(AU\Box)^+$, in this order.

For the ‘if’ direction of rule $(EU\sigma\Box)^+$, let us suppose that $\mathcal{K} \models \Sigma, E\Pi$, where Π contains at least one eventuality. There exists $x \in \text{fullpaths}(\mathcal{K})$ such that $\mathcal{K}, x, 0 \models \Sigma, \Pi$. We are going to prove that there exists \mathcal{K}' such that one of the following properties holds:

- (a) $\mathcal{K}' \models \Sigma, \sigma_2, E(\Diamond\sigma_3, \Pi^{-i})$ for some $\pi_i = \sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3)$ in $\Pi_{\mathcal{U}}$.
- (b) $\mathcal{K}' \models \Sigma, E(\Box\sigma_2, \Pi^{-i})$ for some $\pi_i = \sigma_1 \mathcal{U} \Box\sigma_2$ in $\Pi_{\mathcal{U}}$.
- (c) $\mathcal{K}' \models \Sigma, \sigma_1, E(\circ((\sigma_1 \wedge \neg\Sigma)\mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3)), \Pi^{-i})$ for some $\pi_i = \sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3)$ in $\Pi_{\mathcal{U}}$.
- (d) $\mathcal{K}' \models \Sigma, \sigma_1, E(\circ((\sigma_1 \wedge \neg\Sigma)\mathcal{U}\Box\sigma_2), \Pi^{-i})$ for some $\pi_i = \sigma_1 \mathcal{U} \Box\sigma_2$ in $\Pi_{\mathcal{U}}$.

Since $\mathcal{K}, x, 0 \models \pi_i$ for all $\pi \in \Pi_{\mathcal{U}}$, we define j to be the least $i \geq 0$ such that $\mathcal{K}, x, i \models \varphi$ for some $\sigma_1 \mathcal{U} \varphi \in \Pi_{\mathcal{U}}$. If $j = 0$, then (a) and (b) (depending on the form of φ) are trivially satisfied for $\mathcal{K}' = \mathcal{K}$. Otherwise, if $j > 0$, it holds that $\mathcal{K}, x, j \models \varphi$ and for all $i < j$: $\mathcal{K}, x, i \models \sigma_1$. Moreover, as j is the minimal index, for all $0 \leq i < j$: $\mathcal{K}, x, i \models \sigma'_1$ for all $\sigma'_1 \mathcal{U} \varphi' \in \Pi_{\mathcal{U}}$. Consider k to be the greatest index i such that $0 \leq i < j$ and $\mathcal{K}, x, i \models \Sigma$. Henceforth, we have that $\mathcal{K}, x, k \models \Sigma$ and $\mathcal{K}, x, h \models \neg\Sigma$, for all h such that $k + 1 \leq h < j$. Therefore, (c) and (d) hold for $\mathcal{K}' = \mathcal{K} \upharpoonright x(k)$.

For the ‘if’ direction of the rule $(AU\sigma)^+$, let us suppose that the three sets $\Sigma \cup S_{\Sigma, \beta_1}$, $\Sigma \cup S_{\Sigma, \beta_2}$, and $\Sigma \cup S_{\Sigma, \beta_3}$ of the rule $(AU\sigma)^+$ are unsatisfiable. We will show that the set $\Sigma, A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3), \Pi)$ must be also unsatisfiable. By the hypothesis, we know that any model of Σ is not a model of S_{Σ, β_i} for all $i \in \{1, 2, 3\}$. In other words, for any \mathcal{K} such that $\mathcal{K} \models \Sigma$, the followings three facts holds:

- (a) $\mathcal{K} \not\models \sigma_2 \wedge A(\Diamond\sigma_3, \Pi)$.
- (b) $\mathcal{K} \not\models \sigma_1 \wedge A(\circ((\sigma_1 \wedge (\neg\Sigma \vee \varphi_{\Pi}))\mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3)), \Pi)$.
- (c) $\mathcal{K} \not\models A\Pi$.

To show that $\Sigma, A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3), \Pi)$ is unsatisfiable, we consider an arbitrary \mathcal{K} such that $\mathcal{K} \models \Sigma$ and prove that $\mathcal{K} \not\models A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3), \Pi)$. Since $\mathcal{K} \models \Sigma$, then (a), (b) and (c) hold. According to (b), there are two possible cases:

(Case 1): If $\mathcal{K} \not\models \sigma_1$ then, by (a), either $\mathcal{K} \models \neg\sigma_1 \wedge \neg\sigma_2$ or $\mathcal{K} \models \neg\sigma_1 \wedge E(\Box\neg\sigma_3, \neg\Pi)$. In both cases, it is easy to see that $\mathcal{K} \not\models A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3), \Pi)$.

(Case 2): Otherwise, if $\mathcal{K} \not\models A(\circ((\sigma_1 \wedge (\neg\Sigma \vee \varphi_{\Pi}))\mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3)), \Pi)$, then there exists $x_1 \in \text{fullpaths}(\mathcal{K})$ such that $\mathcal{K}, x_1, 0 \models \circ((\sigma_1 \wedge (\neg\Sigma \vee \varphi_{\Pi}))\mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3)) \wedge \neg\Pi$. This yields two possible cases:

(Case 2.1): If $\mathcal{K}, x_1, 0 \models \circ(\Box\neg\sigma_2 \vee \Box\neg\sigma_3) \wedge \neg\Pi$, then it is trivial that $\mathcal{K} \not\models A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3), \Pi)$.

(Case 2.2): Otherwise, there should exist $i_1 > 0$ that satisfies the following three properties:

- (i) $\mathcal{K}, x_1, j \models (\neg\sigma_2) \vee \Box\neg\sigma_3$ for all j such that $0 \leq j \leq i_1$, and
- (ii) $\mathcal{K}, x_1, i_1 \models \neg\sigma_1 \vee (\Sigma \wedge \neg\varphi_{\Pi})$, and
- (iii) $\mathcal{K}, x_1, 0 \models \neg\Pi$.

If (i) is satisfied because $\mathcal{K}, x_1, j \models \Box\neg\sigma_3$ (for some j such that $0 \leq j \leq i_1$) then trivially $\mathcal{K}, x_1, 0 \not\models \sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3)$. This, along with the fact (iii), not only ensures that $\mathcal{K} \not\models A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3), \Pi)$ but also applies to any other formula $\sigma'_1 \mathcal{U} (\sigma'_2 \wedge \Diamond\sigma'_3)$ in Π . Henceforth, in what follows, we can suppose that for all j such that $0 \leq j \leq i_1$: $\mathcal{K}, x_1, j \models \neg\sigma_2$ and also that $\mathcal{K}, x_1, j \models \neg\sigma'_2$ for all $\sigma'_1 \mathcal{U} (\sigma'_2 \wedge \Diamond\sigma'_3) \in \Pi$.

If (ii) is satisfied because $\mathcal{K}, x_1, i_1 \models \neg\sigma_1$ then it is clear that $\mathcal{K}, x_1, 0 \not\models \sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3)$. Therefore, by (i) and (iii), $\mathcal{K} \not\models A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3), \Pi)$.

Otherwise, if (ii) is satisfied because $\mathcal{K}, x_1, i_1 \models \Sigma \wedge \neg\varphi_{\Pi}$, then (a), (b) and (c) hold for $\mathcal{K} \upharpoonright x_1(i_1)$ (instead of \mathcal{K}) because $\mathcal{K}, x_1, i_1 \models \Sigma$. Hence, applying the same reasoning for $\mathcal{K} \upharpoonright x_1(i_1)$ as we did above for \mathcal{K} , we conclude that there should be a path $x_2 \in \text{fullpaths}(\mathcal{K} \upharpoonright x_1(i_1))$ such that one of the following two facts holds:

(Case 2.2.1): $\mathcal{K} \upharpoonright x_1(i_1), x_2 \models \Box\neg(\sigma_2 \wedge \Diamond\sigma_3) \wedge \neg\Pi$, and therefore $\mathcal{K} \not\models A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3), \Pi)$.

(Case 2.2.2): there should exist $i_2 > 0$ such that $\mathcal{K} \upharpoonright x_1(i_1), x_2, i_2 \models \Sigma \wedge \neg\varphi_{\Pi}$ and for all $j \in \{0..i_2\}$:

- $\mathcal{K} \upharpoonright x_1(i_1), x_2, j \models \neg\sigma_2$, and
- $\mathcal{K} \upharpoonright x_1(i_1), x_2, j \models \neg\sigma'_2$ for all $\sigma'_1 \mathcal{U} (\sigma'_2 \wedge \Diamond\sigma'_3) \in \Pi$

Now, (a), (b) and (c) apply to $\mathcal{K} \upharpoonright x_2(i_2)$. Hence, the infinite iteration of the second case yields a path $y = x_1^{\leq i_1} x_2^{\leq i_2} \dots x_k^{\leq i_k} \dots$ (that exists by the limit closure property) for which the Proposition 32 ensures that $\mathcal{K}, y, 0 \not\models A(\sigma_1 \mathcal{U} (\sigma_2 \wedge \Diamond\sigma_3), \Pi)$.

The proof for the $(AU\Box)^+$ rule follows the same scheme. \square

Theorem 34 (Soundness of the tableau method). *Given any set of state formulae Σ , if there exists a closed tableau for Σ then $\text{UnSat}(\Sigma)$.*

Proof. Let T_{Σ} be a closed tableau for Σ . The set of formulae labelling at least one leaf in each bunch is inconsistent and therefore unsatisfiable. Then, by Lemma 33, the labelling of the root node, Σ , is unsatisfiable. \square

6. Completeness

In this section, we prove the completeness of the introduced tableau method. First, we define the notions of stage, expanded bunch, and expanded tableau. Then we prove the refutational completeness: every unsatisfiable set of state formulae has a closed tableau. In fact, we are going to prove that, for any set Σ_0 , if the systematic tableau $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, given by Algorithm 1, is open, then Σ_0 is satisfiable. That is, $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ has at least one open bunch that allows us to construct a model of Σ_0 . The final step of proving the completeness of the tableau method is establishing its termination.

6.1. Open bunch model construction

In this subsection, we define a method to associate a Kripke structure to any open bunch of the systematic tableau $\mathcal{A}_{\Sigma_0}^{\text{sys}}$. Later, we prove that this Kripke structure is a model of Σ_0 .

Definition 35 (Stage). Given a branch, b of a tableaux T , a *stage* in T is every maximal subsequence of successive nodes n_i, n_{i+1}, \dots, n_j in b such that $\tau(n_k)$ is not a $(\mathbf{Q}\circ)$ -child of $\tau(n_{k-1})$, for all k such that $i < k \leq j$. We denote by $\text{stages}(b)$ the *sequence of all stages* of b . The successor relation on $\text{stages}(b)$ is induced by the successor relation on b . The labelling function τ is extended to stages as the union of the original τ applied to every node in a stage. \square

We prove that any open bunch of the systematic tableau $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ represents a model of the initial set of formulae Σ_0 .

Definition 36 ($\alpha\beta^+$ -saturated stage). A set of state formulae Ψ is $\alpha\beta^+$ -saturated iff for all $\sigma \in \Psi$:

1. If σ is an α -formula then $S_\sigma \subseteq \Psi$.
2. If σ is a β -formula of range k , but it is not a β^+ -formula, then $S_{\beta_i} \subseteq \Psi$ for some $1 \leq i \leq k$.
3. If σ is a β -formula and also a β^+ -formula of range k then either $S_{\beta_i} \subseteq \Psi$ or $S_{\Sigma, \beta_i}^+ \subseteq \Psi$ for some $1 \leq i \leq k$ and $\Sigma = \tau(n) \setminus \{\sigma\}$ for some $n \in s$.

We say that a stage s in $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ is $\alpha\beta^+$ -saturated iff $\tau(s)$ is $\alpha\beta^+$ -saturated. For a given set Σ of state-formulae, we denote by $\text{Comp}(\Sigma)$ the union of all the minimal sets that contains Σ and are $\alpha\beta^+$ -saturated. \square

Definition 37 (Expanded bunch and tableau). An open branch b is *expanded* if each stage $s \in \text{stages}(b)$ is $\alpha\beta^+$ -saturated and b is *eventuality-covered*. A bunch is expanded if all its open branches are expanded. A *tableau* is expanded if all its open bunches are expanded. \square

Proposition 38. Given any set of state formulae Σ_0 , the systematic tableau $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ is expanded.

Proof. Trivial, by construction. \square

Definition 39 (Open bunch model construction). For any expanded bunch H of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, we define the Kripke-structure $\mathcal{K}_H = (S, R, L)$ such that $S = \bigcup_{b \in H} \text{stages}(b)$ and for any $s \in S$: $L(s) = \{p \mid p \in \tau(n) \cap \text{Prop for some node } n \in s\}$; and R is the relation induced in $\text{stages}(b)$ for each $b \in H$. \square

6.2. Properties of the open branches of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$

In order to prove that \mathcal{K}_H , as defined in Definition 39, is a model for the label of the root of H , we first prove the required auxiliary properties of the systematic construction of the tableau $\mathcal{A}_{\Sigma_0}^{\text{sys}}$. The systematic construction of Algorithm 1 produces uniform sets as expandable nodes. The selection is always made in uniform sets and loop-nodes are also labelled by uniform sets.

Remark 40 (Notation for eventualities and tableau rule application). In what follows, we use χ to represent a formula of one of the two following forms: $(\sigma_2 \wedge \diamond\sigma_3)$ or $\square\sigma_2$. Then, $\sigma_1 \mathcal{U} \chi$ stands for one of the two possible eventualities. We say that the *corresponding β^+ rule* is applied to a selected $\mathbf{A}\Pi$ with some marked eventuality $\pi \in \Pi$, meaning that $(\mathbf{A}\mathcal{U}\sigma)^+$ is applied when π is $\sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$, $(\mathbf{A}\mathcal{U}\square)^+$ is applied when π is $\sigma_1 \mathcal{U}\square\sigma_2$, and $(\mathbf{A}\square\mathcal{U})$ followed by $(\mathbf{A}\mathcal{U}\sigma)^+$ is applied when π is $\square(\sigma_1 \mathcal{U}\sigma_2)$. We say that the *corresponding β^+ rule* is applied to a selected $\mathbf{E}\Pi$ meaning that $(\mathbf{E}\mathcal{U}\sigma\square)^+$ is applied when Π contains at least one $\sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ or at least one $\sigma_1 \mathcal{U}\square\sigma_2$, and otherwise when Π contains at least one $\square(\sigma_1 \mathcal{U}\sigma_2)$, then $(\mathbf{E}\mathcal{U}\sigma\square)^+$ is applied just after $(\mathbf{E}\square\mathcal{U})$ has been applied to every $\square(\sigma_1 \mathcal{U}\sigma_2)$ in Π . For clarity, we consider sets S_{Σ, β_i} and S_{Σ, β_i}^+ used in the tableau rules creating child nodes from left ($i = 1$) to right, where i is the rank of the rule. \square

Definition 41 (Variants). For a given set Π of basic path formulae, we denote by $\text{Variants}(\Pi)$ the collection of all subsets of the sets Π' that are obtained from Π by one simultaneous application of any number (including zero) of individual substitutions of some $\pi \in \Pi$ by π' that satisfies the following rules:

- π is an eventuality $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond \sigma_3)$ and π' is either $\diamond \sigma_3$ or a next-step variant of π .
- π is an eventuality $\sigma_1 \mathcal{U} \square \sigma_2$ and π' is either $\square \sigma_2$ or a next-step variant of π .
- π is $\square (\sigma_1 \vee \square \sigma_2)$ and π' is $\square \sigma_2$. \square

The following two propositions establish general properties of the rule-based decomposition of E-conjunctive and A-disjunctive formulae (respectively) along open branches.

Proposition 42. Let b be an open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, let s_i, s_j ($i < j$) be any pair of consecutive stages in $\text{stages}(b)$ and let $\Sigma \cup \{\text{E}\Pi\}$ be the uniform set labelling the first node of s_i . There exists a (possibly empty and minimal) uniform set $\Pi' \in \text{Variants}(\Pi)$ such that $\text{E}\Pi' \in \tau(s_j)$ and for all $\pi \in \Pi$:

- (a) if $\pi = \sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond \sigma_3)$ then there exists $k \geq i$ such that $\sigma_1 \in \tau(s_h)$ for all $h \in i..k-1$ and
 - (a1) $k < j$ and $\sigma_2 \in \tau(s_k)$ and $\sigma_3 \in \tau(s_{k'})$ for some k' such that $k \leq k' \leq j$, or
 - (a2) $k < j$ and $\sigma_2 \in \tau(s_k)$ and $\diamond \sigma_3 \in \Pi'$, or
 - (a3) $k = j$ and π or some next-step variant of it is in Π' ,
- (b) if $\pi = \sigma_1 \mathcal{U} \square \sigma_2$ then there exists $k \geq i$ such that $\sigma_1 \in \tau(s_h)$ for all $h \in i..k-1$ and
 - (b1) $k < j$ and $\sigma_2 \in \tau(s_h)$ for some h such that $i \leq h \leq k$ and $\square \sigma_2 \in \Pi'$, or
 - (b2) $k = j$ and $\sigma_1 \mathcal{U} \square \sigma_2$ or some next-step variant of it is in Π' ,
- (c) if $\pi = \square (\sigma_1 \mathcal{U} \sigma_2)$ then $\square (\sigma_1 \mathcal{U} \sigma_2) \in \Pi'$ and for all $h \in i..j-1$: $\sigma_2 \in \tau(s_h)$ or $\sigma_1 \in \tau(s_h)$,
- (d) if $\pi = \square (\sigma_1 \vee \square \sigma_2)$ then there exists $k \geq i$ such that $\sigma_1 \in \tau(s_h)$ for all $h \in i..k-1$,
 - (d1) $k = j$ and $\square (\sigma_1 \vee \square \sigma_2) \in \Pi'$, or
 - (d2) $k < j$ and there exists $k' \in k..j-1$ such that $\sigma_2 \in \tau(s_h)$ for all $h \in k'..j-1$ and $\square \sigma_2 \in \Pi'$.

Proof. By simultaneous induction on Π , using the rules for the E-conjunctive formulae. Note that any stage in b is $\alpha\beta^+$ -saturated and the procedure `Uniform_Tableau` applies exactly once between the last state of s_i and the first state of s_{i+1} . More specifically, in (a) we use $(\text{E}\mathcal{U}\sigma)$ and $(\text{E}\mathcal{U}\sigma\square)^+$, and (a1) and (a2) come from the second application of $(\text{E}\mathcal{U}\sigma)$ to $\text{E}(\diamond\sigma_3, \dots)$ where $\diamond\sigma_3$ abbreviates $\tau\mathcal{U}\sigma_3$. Similarly, item (b) comes from rules $(\text{E}\mathcal{U}\square)$ and $(\text{E}\mathcal{U}\sigma\square)^+$; Items (c) and (d) are respectively obtained from the rules $(\text{E}\square\mathcal{U})$, $(\text{E}\mathcal{U}\sigma)$, and $(\text{E}\mathcal{U}\sigma\square)^+$; and $(\text{E}\square\sigma)$. It is worth noting that Π' is empty if all the formulae in Π satisfy the case (a1). \square

Proposition 43. Let b be an open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, let s_i, s_j ($j > i$) be any pair of consecutive stages in $\text{stages}(b)$ and let $\Sigma \cup \{\text{A}\Pi\}$ be the uniform set labelling the first node of s_i . If there exists a non-empty uniform set $\Pi' \in \text{Variants}(\Pi)$ such that $\text{A}\Pi' \in \tau(s_j)$, then the following four facts hold:

- (a) For all $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond \sigma_3) \in \Pi$:
 - (a1) if $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond \sigma_3)$ or a next-step variant of it is in Π' , then $\sigma_1 \in \tau(s_h)$ for all $h \in i..j-1$,
 - (a2) if $\diamond \sigma_3 \in \Pi'$ then $\sigma_2 \in \tau(s_k)$ for some $k \in i..j-1$ and $\sigma_1 \in \tau(s_h)$ for all $h \in i..k-1$.
- (b) For all $\sigma_1 \mathcal{U} \square \sigma_2 \in \Pi$: if $\square \sigma_2 \in \Pi'$ or $\sigma_1 \mathcal{U} \square \sigma_2$ or a next-step variant of $\sigma_1 \mathcal{U} \square \sigma_2$ is in Π' , then $\{\sigma_1, \sigma_2\} \subseteq \tau(s_h)$ for all $h \in i..j-1$.
- (c) For all $\square (\sigma_1 \mathcal{U} \sigma_2) \in \Pi \cap \Pi'$: $\sigma_1 \in \tau(s_h)$ or $\sigma_2 \in \tau(s_h)$ for all $h \in i..j-1$.
- (d) For all $\square (\sigma_1 \vee \square \sigma_2) \in \Pi$,
 - (d1) if $\square (\sigma_1 \vee \square \sigma_2) \in \Pi$ then $\sigma_1 \in \tau(s_h)$ for all $h \in i..j-1$,
 - (d2) if $\square \sigma_2 \in \Pi'$ then for some $k \in i..j-1$ $\{-\sigma_1, \sigma_2\} \subseteq \tau(s_k)$ and $\sigma_1 \in \tau(s_h)$ for all $h \in i..k-1$ and $\sigma_2 \in \tau(s_h)$ for all $h \in k..j-1$.

Moreover, if there exists no $\text{A}\Pi' \in \tau(s_j)$ such that $\Pi' \in \text{Variants}(\Pi)$, then there exists some $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond \sigma_3) \in \Pi$ and some $k, k' \in i..j-1$ such that $k \leq k'$ and $\sigma_2 \in \tau(s_k)$ and $\sigma_3 \in \tau(s_{k'})$.

Proof. By simultaneous induction on Π . For the proof we note that any stage in b is $\alpha\beta^+$ -saturated, and the following rules for A-disjunctive formulae hold: $(\text{A}\mathcal{U}\sigma)$ and $(\text{A}\mathcal{U}\sigma)^+$ in (a); $(\text{A}\mathcal{U}\square)$ and $(\text{A}\mathcal{U}\square)^+$ in (b); $(\text{A}\square\mathcal{U})$, $(\text{A}\mathcal{U}\sigma)$, and $(\text{A}\mathcal{U}\sigma)^+$ in (c); and $(\text{A}\square\sigma)$ in (d). It is worth noting that the rules $(\text{A}\mathcal{U}\sigma)$ and $(\text{A}\square\sigma)$ generate a child where the A-disjunctive formula that comes from $\text{A}\Pi$ is of the form $\text{A}\Pi'$ with $\Pi' \subset \Pi$ (one formula is lost). \square

From the previous two propositions it is easy to conclude that, for a given initial node (of stage s_i) labelled by a uniform set $\Sigma \cup \{\text{Q}\Pi\}$, the mentioned sets Π' such that $\text{Q}\Pi' \in \tau(s_{i+1})$ exclusively consist of path subformulae and next-step variants of the formulas in Π . It also follows that $\tau(s_{i+1})$ exclusively contains Boolean combinations of state subformulae in $\Sigma \cup \{\text{Q}\Pi\}$

and the formula $\mathbf{Q}\Pi'$. The next proposition shows that the selection of an E-conjunctive formula in any open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ can only be kept along finitely many stages.

Proposition 44. *Let b be an open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, let $s_i \in \text{stages}(b)$ and let $\Sigma \cup \{\mathbf{E}\Pi\}$ be the uniform set labelling the first node of s_i where the selectable formula $\mathbf{E}\Pi$ is selected. If $\mathbf{E}\Pi$ has the highest priority, then*

- (a) *For all $\pi = \sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond\sigma_3) \in \Pi: \{\sigma_2, \mathbf{E}(\diamond\sigma_3, \Pi')\} \subseteq \tau(s_k)$ for some stage $s_k \in \text{stages}(b)$ ($k \geq i$) and some $\Pi' \in \text{Variants}(\Pi \setminus \{\pi\})$.*
- (b) *For all $\pi = \sigma_1 \mathcal{U}\square\sigma_2 \in \Pi: \mathbf{E}(\square\sigma_2, \Pi') \in \tau(s_k)$ for some stage $s_k \in \text{stages}(b)$ ($k \geq i$) and some $\Pi' \in \text{Variants}(\Pi \setminus \{\pi\})$,*

and if $\mathbf{E}\Pi$ has the lowest priority, then

- (c) *For all $\pi = \square(\sigma_1 \mathcal{U}\sigma_2) \in \Pi: \sigma_2 \in \tau(s_k)$ for some stage $s_k \in \text{stages}(b)$ ($k \geq i$).*

Proof. Suppose that $\mathbf{E}\Pi$ has the highest priority, then Π contains at least one $\sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ or $\sigma_1 \mathcal{U}\square\sigma_2$, then the β^+ -rule $(\mathbf{E}\mathcal{U}\sigma\square)^+$ is applied in the first node of stage s_i where $\mathbf{E}\Pi$ is selected and the resulting $\mathbf{E}\Pi'$ of the highest priority in its children are kept selected, hence $(\mathbf{E}\mathcal{U}\sigma\square)^+$ is again applied to them, and so on. We proceed by contradiction, supposing that (a) and (b) does not hold. We get a contradiction from the hypothesis that (a) does not hold, i.e. for some $\sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond\sigma_3) \in \Pi: \{\sigma_2, \mathbf{E}(\diamond\sigma_3, \Pi')\} \not\subseteq \tau(s_k)$ for every stage $s_k \in \text{stages}(b)$ such that $k \geq i$ and all $\Pi' \in \text{Variants}(\Pi \setminus \{\pi_i\})$. The proof for the case where (b) does not hold is identical. The fact that (a) does not hold means that at the first node of every stage $s_j \in \text{stages}(b)$ such that $j \geq i$ there is a selected formula $\mathbf{E}\Pi_{s_j}$ which satisfies that $\sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ or some step-variant of it is in Π_{s_j} . Hence, except for a finite number of applications of $(\mathbf{E}\mathcal{U}\sigma\square)^+$ that extends the branch b with some set $S_{\Sigma_{s_j}, \beta_i}^+$ (where Σ_{s_j} is the context of the selected formula at the first node of each stage s_j , in particular $\Sigma_{s_i} = \Sigma$) such that $1 \leq i \leq n$, the branch b is repeatedly extended with some set $S_{\Sigma_{s_j}, \beta_i}^+$ such that $n+1 \leq i \leq 2n$, which includes a next-step variant of at least one formula in Π_{s_j} . Note that this next-step variant could be of $\sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$ or some other formula of the form $\sigma \mathcal{U}(\sigma \wedge \diamond\sigma)$ or $\sigma \mathcal{U}\square\sigma$. In any case, the uniform set labelling the first node of each stage s_j ($i \leq j$) has the form $\Sigma_{s_j}, \mathbf{E}\Pi_{s_j}$ where Π_{s_j} contains at least one next-step variant of $\sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$, $\mathbf{E}\Pi_{s_j}$ is the selected formula and for every (except a finite number of) Σ_{s_k} where $i \leq k \leq j$, Π_{s_j} contains a formula of the form $\sigma \wedge (\neg\Phi_1 \wedge \dots \wedge \neg\Phi_r) \mathcal{U}\chi$ such that $\Phi_h = \Sigma_{s_k}$ for some $1 \leq h \leq r$. Since no other β^+ -rule is applied each Σ_{s_j} is a subset of the finite set formed by all state formulae that are subformulae of some formula in $\Sigma_{s_i} \cup \Pi$ and their negations. Hence, there exists a finite number of different Σ_{s_j} . Therefore, after finitely many applications of the β^+ -rule $(\mathbf{E}\mathcal{U}\sigma\square)^+$, for some $k > i$, $\Sigma_{s_k} = \Sigma_{s_h}$ for some $h \in \{i, \dots, k-1\}$, and $\neg\Sigma_{s_k} \in \tau(s_k)$, hence, Σ_{s_k} must be inconsistent. Since b is open, this is a contradiction. It means that for some $k \geq i$, the application of the corresponding β^+ -rule $(\mathbf{E}\mathcal{U}\sigma\square)^+$ should produce a node whose label contains $S_{\Sigma_{s_j}, \beta_i}^+$ where $1 \leq i \leq n$. Henceforth, the open branch b must satisfy (a).

If $\mathbf{E}\Pi$ has the lowest priority, then Π contains at least one $\square(\sigma \mathcal{U}\sigma)$ (and none $\sigma \mathcal{U}(\sigma \wedge \diamond\sigma)$ and none $\sigma \mathcal{U}\square\sigma$). Let us suppose there are $n \geq 1$ formulae, i.e. $\mathbf{E}\Pi = \mathbf{E}(\square(\sigma_1^1 \mathcal{U}\sigma_2^1), \dots, \square(\sigma_1^n \mathcal{U}\sigma_2^n), \Pi')$ for some Π' that does not contain any eventuality. Then, the α -rule $(\mathbf{E}\square\mathcal{U})$ is applied n times transforming the selected $\mathbf{E}\Pi$ into

$$\mathbf{E}(\sigma_1^1 \mathcal{U}\sigma_2^1, \dots, \sigma_1^n \mathcal{U}\sigma_2^n, \circ(\sigma_1^1 \mathcal{U}\sigma_2^1), \dots, \circ(\sigma_1^n \mathcal{U}\sigma_2^n), \Pi')$$

which has the highest priority. Hence, by a particular application of the case (a), for all $1 \leq j \leq n: \sigma_2^j \in \tau(s_k)$ for some stage $s_k \in \text{stages}(b)$ ($k \geq i$). \square

For A-disjunctive formulae, not only the outer context, but also the inner context, plays an important role. The next propositions explain the role of both kinds of contexts.

Proposition 45. *Let b be an open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, let $s_i \in \text{stages}(b)$ and let $\Sigma_{s_i} \cup \{\mathbf{A}\Pi\}$ be the uniform set labelling the first node of s_i where $\mathbf{A}\Pi$ is selected and some eventuality $\pi_{\mathcal{U}} \in \Pi$ is marked. Let b be any branch where the next-steps variants of $\pi_{\mathcal{U}}$ are successively marked and $\Pi_{s_i} = \Pi \setminus \{\pi_{\mathcal{U}}\}$ is the initial inner context. Then, one of the following two facts hold:*

- (a) *There exists $k \geq i$ and some $\Pi' \in \text{Variants}(\Pi \setminus \{\pi_{\mathcal{U}}\})$ such that:*
 - (a1) $\{\sigma_2, \mathbf{A}(\diamond\sigma_3, \Pi')\} \subseteq \tau(s_k)$ if $\pi_{\mathcal{U}} = \sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond\sigma_3)$.
 - (a2) $\mathbf{A}(\square\sigma_2, \Pi') \in \tau(s_k)$ if $\pi_{\mathcal{U}} = \sigma_1 \mathcal{U}\square\sigma_2 \in \Pi'$.
 - (a3) $\sigma_2 \in \tau(s_k)$ if $\pi_{\mathcal{U}} = \square(\sigma_1 \mathcal{U}\sigma_2) \in \Pi'$.
- (b) *There exists $k \geq i$ such that the first node of s_k is a loop-node whose companion node is in s_h , for some $h \in \{i..k-1\}$, some $\Pi' \in \text{Variants}(\Pi \setminus \{\pi_{\mathcal{U}}\})$ and some next-step variant $\pi_{\mathcal{U}}^v$ of $\pi_{\mathcal{U}}$ such that $\mathbf{A}(\pi_{\mathcal{U}}^v, \Pi') \in \tau(s_j)$ for all $j \in \{h..k\}$, and $\varphi_{\Pi'} \in \tau(s_k)$.*

Moreover, in both cases, for all $j \in \{i..k-1\}$:

1. $\sigma_1 \in \tau(s_j)$ for all $\sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond \sigma_3) \in \Pi' \cup \{\pi_{\mathcal{U}}\}$,
2. $\{\sigma_1, \sigma_2\} \subseteq \tau(s_j)$ for all $\sigma_1 \mathcal{U} \square \sigma_2 \in \Pi' \cup \{\pi_{\mathcal{U}}\}$,
3. $\sigma_1 \in \tau(s_j)$ or $\sigma_2 \in \tau(s_j)$ for all $\square(\sigma_1 \mathcal{U} \sigma_2) \in \Pi' \cup \{\pi_{\mathcal{U}}\}$, and
4. $\sigma_1 \in \tau(s_j)$ for all $\square(\sigma_1 \vee \square \sigma_2) \in \Pi'$.

Proof. If $\varphi_{\pi_{s_i}} = \mathbf{F}$, the uniform set labelling the first node at each stage s_j ($j \geq i$) of b has the form

$$\Sigma_{s_j}, \mathbf{A}((\sigma_1 \wedge (\neg \Sigma_{s_i} \wedge \dots \wedge \neg \Sigma_{s_{j-1}})) \mathcal{U} \chi, \Pi_{s_j})$$

where each Σ_{s_j} is the outer (resp. inner) context of the selected formula containing the marked next-step variant of $\pi_{\mathcal{U}}$ at the first node of each stage s_j . In particular, $\Sigma_{s_i} = \Sigma$. Since no other β^+ -rule is applied each Σ_{s_j} is a subset of the finite set formed by all state formulae that are subformulae of some formula in $\Sigma_{s_i} \cup \Pi$ and their negations. Hence, there are a finite number of different Σ_{s_j} . Therefore, after finitely many applications of the β^+ -rule, $\Sigma_{s_h} = \Sigma_{s_j}$, for some $h \geq i$, for some $j \in \{i, \dots, h-1\}$, and $\sigma_1 \wedge (\neg \Sigma_{s_i} \wedge \dots \wedge \neg \Sigma_{s_{h-1}}) \in \tau(s_h)$. In particular, $\neg \Sigma_{s_h} \in \tau(s_h)$, hence, Σ_{s_h} must be inconsistent. Since b is open, this is a contradiction. This means that, for some $k \geq i$ the application of the corresponding β^+ -rule should produce a node generated by the corresponding set $S_{\beta_1}^+$. Henceforth, the open branch b must satisfy (a1) or (a2) or (a3), depending on the case of $\pi_{\mathcal{U}}$.

If $\varphi_{\pi_{s_i}} \neq \mathbf{F}$, then, the uniform set labelling the first node at each stage s_j ($j \geq i$) has the form

$$\Sigma_{s_j}, \mathbf{A}((\sigma_1 \wedge (\neg \Sigma_{s_i} \vee \varphi_{\Pi_{s_i}}) \wedge \dots \wedge (\neg \Sigma_{s_{j-1}} \vee \varphi_{\Pi_{s_{j-1}}})) \mathcal{U} \chi, \Pi_{s_j})$$

where each Π_{s_h} is the inner context at the first node of each stage s_h . In particular, $\Pi_{s_i} = \Pi \setminus \{\pi_{\mathcal{U}}\}$. Since no other β^+ -rule is applied, then

- each Σ_{s_j} is a subset of the following finite set $LC(\Sigma, \Pi)$: $LC(\Sigma, \Pi)$ is formed by all state formulae that are subformulae of some formula in $\Sigma \cup \Pi$, their negations, and a formula $E \diamond \sigma_2$ for each subformula $\square(\sigma_1 \mathcal{U} \sigma_2)$ in Π (see Definition 19), and
- each Π_{s_j} is a subset of the finite set of all state formulae that are subformulae of some formula in Π . Indeed, each $\Pi_{s_{j+1}} \in \text{Variants}(\Pi_{s_j})$ for all $j \geq i$.

In particular, there are a finite number of different outer and inner contexts. Henceforth, there are two possibilities. First, for some h, k such that $k > h \geq i$, both $\Sigma_{s_k} = \Sigma_{s_h}$ and $\Pi_{s_k} = \Pi_{s_h}$; and second for some $h \geq i$, the formula $\varphi_{\pi_{s_h}}$ is \mathbf{F} . In the latter case, the item (a) must be satisfied for some $k \geq h$. In the former case, by Definition 41 and Proposition 43, for all $j \in \{h..k\}$: $\Pi_{s_j} = \Pi_{s_k}$. Let $\Pi' = \Pi_{s_k}$, then the first nodes at the sequence of stages s_h, s_{h+1}, \dots, s_k are respectively labelled by

$$\Sigma_{s_h} \cup \{\mathbf{A}((\sigma_1 \wedge \delta) \mathcal{U} \chi, \Pi')\}, \Sigma_{s_{h+1}} \cup \{\mathbf{A}((\sigma_1 \wedge \delta) \mathcal{U} \chi, \Pi')\}, \dots, \Sigma_{s_k} \cup \{\mathbf{A}((\sigma_1 \wedge \delta) \mathcal{U} \chi, \Pi')\}$$

where $\delta = (\neg \Sigma_{s_i} \vee \varphi_{\Pi_{s_i}}) \wedge \dots \wedge (\neg \Sigma_{s_h} \vee \varphi_{\Pi'}) \wedge \dots \wedge (\neg \Sigma_{s_{k-1}} \vee \varphi_{\Pi'})$ or equivalently $\delta = (\neg \Sigma_{s_i} \vee \varphi_{\Pi_{s_i}}) \wedge \dots \wedge ((\neg \Sigma_{s_h} \wedge \dots \wedge \neg \Sigma_{s_{k-1}}) \vee \varphi_{\Pi'})$. Hence, in node s_k , the application of the β^+ -rule to the marked eventuality produces a right-hand child that contains Σ_{s_k} and $\sigma_1 \wedge \delta$. Therefore, by rules (\wedge) and (\vee), it also contains $\neg \Sigma_{s_h} = \neg \Sigma_{s_k}$. Therefore, since b is open, $\tau(s_k)$ must contain $\varphi_{\Pi'}$, which completes the proof of item (b). Moreover, in both cases (a) and (b), for all $j \in \{i..k-1\}$, each inner context $\Pi_{s_{j+1}}$ satisfies the properties of Π' in Proposition 42 with respect to Π_{s_j} as Π . Consequently, the last four items of the proposition hold. \square

The next two propositions provide a detailed description of how the highest priority formulae evolve in open branches.

Proposition 46. Let b be an open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, and let $E \Pi$ be of the highest priority that is selected at some stage $s_i \in \text{stages}(b)$. Then there exists a state $s_k \in \text{stages}(b)$ (for some $k \geq i$) and some (possibly empty and minimal) set $\Pi' \in \text{Variants}(\Pi)$ such that $E \Pi' \in \tau(s_k)$, $E \Pi'$ is of the lowest priority and for all $\pi \in \Pi$ the following facts hold:

- (a) If $\pi = \sigma_1 \mathcal{U}(\sigma_2 \wedge \diamond \sigma_3)$ then there exists j, j' such that $i \leq j \leq j' \leq k$, for all $h \in \{i, \dots, j-1\}$: $\sigma_1 \in \tau(s_h)$, $\sigma_2 \in \tau(s_j)$, and $\sigma_3 \in \tau(s_{j'})$.
- (b) If $\pi = \sigma_1 \mathcal{U} \square \sigma_2$ then there exists j such that $i \leq j \leq k$ and for all $h \in \{i, \dots, j-1\}$: $\sigma_1 \in \tau(s_h)$, and for all $h \in \{j, \dots, k\}$: $\sigma_2 \in \tau(s_h)$ and $\square \sigma_2 \in \Pi'$.
- (c) If $\pi = \square(\sigma_1 \mathcal{U} \sigma_2)$ then $\square(\sigma_1 \mathcal{U} \sigma_2) \in \Pi'$, and for all $j \in \{i, \dots, k\}$: either $\sigma_1 \in \tau(s_j)$ or $\sigma_2 \in \tau(s_j)$.
- (d) If $\pi = \square(\sigma_1 \vee \square \sigma_2)$ then one of the following two facts holds:
 - (d1) For all $j \in \{i, \dots, k\}$: $\sigma_1 \in \tau(s_j)$ and $\square(\sigma_1 \vee \square \sigma_2) \in \Pi'$.
 - (d2) There exists j such that $i \leq j \leq k$ and for all $h \in \{i, \dots, j-1\}$: $\sigma_1 \in \tau(s_h)$, and for all $h \in \{j, \dots, k\}$: $\sigma_2 \in \tau(s_h)$ and $\square \sigma_2 \in \Pi'$.

Proof. By simultaneous induction on the structures of formulae in Π and Propositions 44 and 42((a) and (b)), the above items (a) and (b) hold for some $k \geq i$. Note that in case (a), $E(\diamond\sigma_3, \Pi')$ (for some Π' such that $\{\diamond\sigma_3\} \cup \Pi' \in \text{Variants}(\Pi)$) is kept selected at stage s_j . Hence, the eventuality $\sigma_3 \in \tau(s_{j'})$ for some $j' \geq j$. Therefore, by Proposition 44, the existence of such j' is ensured. In case (b) $E(\Box\sigma_2, \Pi') \in \tau(s_j)$ for some $j \geq i$, hence, by Proposition 42(d), for all $h \in \{j, \dots, k\}$: $\sigma_2 \in \tau(s_h)$ and $\Box\sigma_2 \in \Pi'$. Items (c) and (d) are ensured by Propositions 42 (c) and (d). Additionally, by minimality, Π' only contains formulae of the forms $\Box(\sigma_1 \mathcal{U} \sigma_2)$ and $\Box(\sigma_1 \vee \Box\sigma_2)$, hence $E\Pi'$ is of the priority. \square

The other kind of the highest-priority formulae are $A\Pi$ such that Π is exclusively formed by formulae of the form $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond\sigma_3)$.

Proposition 47. *Let b be an open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, and let $A\Pi$ be a formula of the highest priority that is selected at some stage $s_i \in \text{stages}(b)$. Then there exists $\pi = \sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond\sigma_3) \in \Pi$ and some stage $s_k \in \text{stages}(b)$ (for some $k \geq i$) such that for all $j \in \{i, \dots, k-1\}$: $\sigma_1 \in \tau(s_j)$, $\{\sigma_2, A(\diamond\sigma_3, \Pi')\} \subseteq \tau(s_k)$ for some $\Pi' \in \text{Variants}(\Pi \setminus \{\pi\})$. Moreover, $A(\diamond\sigma_3, \Pi')$ is also a formula of the highest priority.*

Proof. According to Definitions 24 and 26, one eventuality in Π must be marked at the stage s_i of b . Hence, there exists $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond\sigma_3) \in \Pi$ that is the marked eventuality at stage s_i . Since φ_Π is \mathbb{F} when Π is exclusively formed by formulae of the form $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond\sigma_3)$, the item (a) of Proposition 45 holds. Hence, by Proposition 45 (a1), there exists $k \geq i$ and Π' such that $\{\sigma_2, A(\diamond\sigma_3, \Pi')\} \in \tau(s_k)$. Since $\Pi' \in \text{Variants}(\Pi \setminus \{\pi\})$, every formula in Π' is of the form $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond\sigma_3)$ (in particular, $\diamond\sigma$ which abbreviates $\tau \mathcal{U} (\sigma \wedge \diamond\tau)$). Hence, $A(\diamond\sigma_3, \Pi')$ is also of the highest-priority. \square

Next, we show that any open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ is eventuality-covered. In the sequel, we deal with uniform sets formed by non-selectable and the lowest priority formulae (i.e., without the highest-priority formulae), we call them *cycle-uniform* sets.

Proposition 48. *Any open branch b of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ is eventuality-covered.*

Proof. Let b be any open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, we are going to show that there must exist some stage s_ℓ in b with the first node n_ℓ labelled by a cycle-uniform set Σ_ℓ such that any selection of a formula of the lowest priority in Σ_ℓ produces a loop-node whose companion node is n_ℓ .

Let Σ and Σ' be any two cycle-uniform sets of state formulae, we say that $\Sigma' \leq \Sigma$ iff every formula in Σ' is either a proper subformula of some formula $Q\Pi \in \Sigma$ or its negation, or a formula $Q\Pi'$ such that there exists $Q\Pi \in \Sigma$ such that $\Pi' \in \text{Variants}(\Pi)$. It is worth noting that the formulae $E\diamond\sigma_2$ that can be introduced by φ_Π (see Definition 19) are of the highest priority, then they cannot belong to any cycle-uniform set. Let $b = s_0, \dots, s_i, \dots, s_j, \dots$ ($0 \geq i > j$) be any open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, and let Σ and Σ' respectively be the cycle-uniform sets labelling the first node of s_i and s_j . By Propositions 42, 43, 44 and 45, $\Sigma' \leq \Sigma$. Moreover, for any cycle-uniform set Σ , \leq is a well-founded order on the collection of all cycle-uniform sets Σ' such that $\Sigma' \leq \Sigma$.

Let $b = s_0, \dots, s_i, \dots, s_j, \dots, s_k, \dots$ ($0 \geq i > j \geq k$) be any open branch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$. Suppose that every highest priority formula in the initial uniform set has been selected before the stage s_i . Let Σ be the cycle-uniform set labelling the first node of s_j which is a loop-node whose companion is the first node of s_i . Suppose that Σ contains at least one lowest priority formula $A\Pi$ that was selected at s_i . If b is not already eventuality-covered, this means that there exists $A\Pi' \in \Sigma$ of the lowest priority that has not been selected. Suppose that $A\Pi'$ is selected at s_j and there exists a loop-node at s_k labelled by Σ' then, $\Sigma' \leq \Sigma$. If $\Sigma = \Sigma'$ and there are no more selectable formula in Σ , b is already eventuality covered and the first node of s_i is n_ℓ . Otherwise, $\Sigma' < \Sigma$, so that the companion node of the first node of s_k is the first node of some stage s_h such that $h > i$. In general, for any number of the lowest priority formulae in Σ , by well-foundedness of $<$ there should exist a minimal node n_ℓ labelled by a cycle-uniform set Σ_ℓ such that any selection of the lowest priority formula in Σ_ℓ produces a loop-node whose companion node is n_ℓ . Hence, the branch b ends by a subsequence of $n \geq 2$ (possibly non-consecutive) stages s_{i_1}, \dots, s_{i_n} whose first node is labelled by Σ_ℓ , where n_ℓ is the first node of s_{i_1} , and each selectable (lowest priority) formula in Σ_ℓ is selected at some stage in s_{i_1}, \dots, s_{i_n} . In particular, Σ_ℓ could be empty, then $n = 2$ and b is trivially eventuality-covered. \square

6.3. Refutational completeness

In this subsection we prove that our tableau method is refutationally complete, that is if a set of state formulae is unsatisfiable then there exists a closed tableau for it. For that, we first ensure the existence of a model for any open bunch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$.

Lemma 49 (Model existence). *Let H be an expanded bunch of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ and $\mathcal{K}_H = (S, R, L)$ be as in Definition 39. For every state $s \in S$, if $\sigma \in L(s)$ then $\mathcal{K}_H, s, 0 \models \sigma$. Therefore, $\mathcal{K}_H \models \Sigma$.*

Proof. Let H be any expanded bunch of $\mathcal{A}_{\Sigma_0}^{sys}$ and let b be any open branch in H . The construction of any branch of $\mathcal{A}_{\Sigma_0}^{sys}$ starts by selecting a formula of the highest-priority (if any) and marking eventualities as explained in Definition 24. At most one eventuality is marked inside the unique selected QP and the rules of Fig. 7 are applied to this formula and to no one else. When a β^+ -rule is applied to a formula of the highest priority (independently of the marked eventuality), then only the outer context (but no the inner context) is used to construct the new-step variant. Therefore, while some highest priority formulae is selected, previous labels cannot be repeated. Consequently, the initial segment of any open branch has no loop-nodes. This initial segment can be empty or not. According to Proposition 48, the branch b is eventuality covered. Hence, there exists a (possibly empty) cycle-uniform set Σ_ℓ such that for some $i \geq 0$: $b = s_0, s_1, \dots, s_{i-1}, s_i, s_{i+1}, \dots, s_j, n_\ell$, where each s_h stands for a stage and n_ℓ is a non-expandable loop-node labelled by Σ_ℓ whose companion node is the first node at stage s_i . Let $s_{i_1}, s_{i_2}, \dots, s_{i_z}$ be the subsequence formed by all the stages in s_i, s_{i+1}, \dots, s_j whose first node is labelled by Σ_ℓ (in particular, $s_{i_1} = s_i$). Then, each lowest priority formula in Σ_ℓ has been selected at some node n_h ($h \in i..j+1$). The tableau branch b represents a cyclic branch (of a model) such that $\text{path}(b) = s_0, s_1, \dots, s_{i-1} \langle s_i, s_{i+1}, \dots, s_j \rangle^\omega$ where each state s_h ($h \in 0..j$) is labelled by the set of all atoms occurring in the label $\tau(s_h)$ of the tableau stage s_h (in the label of some tableau node at stage s_h).

We are going to prove that $\mathcal{K}_{H, s_a, 0} \models \sigma$ for any $a \in 0..j$ and any formula σ in $\tau(s_a)$, by structural induction on the formula σ . The base of the induction $\sigma = p \in \text{Prop}$ is ensured by Definition 39: $\mathcal{K}_{H, s_a, 0} \models p$.

The bunch H allows us to ensure that whenever a tableau node in stage s_a is labelled by an elementary set $\{\Sigma, A\circ\Phi_1, \dots, A\circ\Phi_n, E\circ\Psi_1, \dots, E\circ\Psi_m\} \subseteq L(s)$ then, by rule (Q \circ), the bunch H contains one successor stage s_{a+1}^i , for each $i \in 1..m$, that contains $\{A\Phi_1, \dots, A\Phi_n, E\Psi_i\}$. Since by induction hypothesis, we can assume that $\mathcal{K}_{H, s_{a+1}^i, 0} \models A\Phi_1, \dots, A\Phi_n, E\Psi_i$ for all $i \in 1..m$, and Σ is a consistent set of literals, then we can infer that $\mathcal{K}_{H, s_a, 0} \models \{\Sigma, A\circ\Phi_1, \dots, A\circ\Phi_n, E\circ\Psi_1, \dots, E\circ\Psi_m\}$.

To complete the proof, we prove different cases for σ being a formula of the form QP, depending on whether σ is selectable or not and, in the selectable case, depending on the σ priority for the selection strategy.

Let $\sigma = E\Pi \in \tau(s_a)$ be non-selectable. Then every $\pi \in \Pi$ is of the form $\Box(\sigma_1 \vee \Box\sigma_2)$. Hence, by Proposition 42 (d) and the induction hypothesis, there exists a state s_k (for some $k \geq a$) and some non-empty set $\Pi' \in \text{Variants}(\Pi)$ such that $\mathcal{K}_{H, s_k, 0} \models E\Pi'$, $\Box(\sigma_1 \vee \Box\sigma_2) \in \Pi'$, and for all $\pi \in \Pi$:

- $\mathcal{K}_{H, s_j, 0} \models \sigma_1$ for all $a \leq j \leq k$, and
- there exists j such that $a \leq j \leq k$ and for all $h \in a..j-1$: $\mathcal{K}_{H, s_h, 0} \models \sigma_1$, and for all $h \in j..k$: $\mathcal{K}_{H, s_h, 0} \models \sigma_2$ and $\Box\sigma_2 \in \Pi'$.

Therefore, $\mathcal{K}_{H, s_a, 0} \models E\Pi$.

Let $\sigma = A\Pi \in \tau(s_a)$ be non-selectable. Then every $\pi \in \Pi$ is of the form $\Box(\sigma_1 \vee \Box\sigma_2)$. By Proposition 43 (d) and the induction hypothesis, there exists a state s_k (for some $k \geq a$) and some non-empty set $\Pi' \in \text{Variants}(\Pi)$ such that $\mathcal{K}_{H, s_k, 0} \models A\Pi'$ and for all $\pi = \Box(\sigma_1 \vee \Box\sigma_2) \in \Pi$:

- If $\Box(\sigma_1 \vee \Box\sigma_2) \in \Pi'$ then $\mathcal{K}_{H, s_j, 0} \models \sigma_1$ for all $j \in a..k$, and
- if $\Box\sigma_2 \in \Pi'$ then there exists j such that $a \leq j \leq k$ and for all $h \in a..j-1$: $\mathcal{K}_{H, s_h, 0} \models \sigma_1$, and for all $h \in j..k$: $\mathcal{K}_{H, s_h, 0} \models \sigma_2$.

Therefore, $\mathcal{K}_{H, s_a, 0} \models A\Pi$.

Let $\sigma = E\Pi \in \tau(s_a)$ be a (selectable) formula of the highest priority. According to Proposition 46 and the induction hypothesis, there exists a state s_k (for some $k \geq a$) and some (possibly empty and minimal) set $\Pi' \in \text{Variants}(\Pi)$ such that $\mathcal{K}_{H, s_k, 0} \models E\Pi'$ and for all $\pi \in \Pi$ the following facts hold:

- If $\pi = \sigma_1 \mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3)$ then there exists j, j' such that $a \leq j \leq j' \leq k$, for all $h \in i..j-1$: $\mathcal{K}_{H, s_h, 0} \models \sigma_1$, $\mathcal{K}_{H, s_j, 0} \models \sigma_2$, and $\mathcal{K}_{H, s_{j'}, 0} \models \sigma_3$.
- If $\pi = \sigma_1 \mathcal{U}\Box\sigma_2$ then there exists j such that $a \leq j \leq k$ and for all $h \in a..j-1$: $\mathcal{K}_{H, s_h, 0} \models \sigma_1$, and for all $h \in j..k$: $\mathcal{K}_{H, s_h, 0} \models \sigma_2$ and $\Box\sigma_2 \in \Pi'$.
- If $\pi = \Box(\sigma_1 \mathcal{U}\sigma_2)$ then $\Box(\sigma_1 \mathcal{U}\sigma_2) \in \Pi'$, and for all $j \in a..k$: either $\mathcal{K}_{H, s_j, 0} \models \sigma_1$ or $\mathcal{K}_{H, s_a, 0} \models \sigma_2$.
- If $\pi = \Box(\sigma_1 \vee \Box\sigma_2)$ then one of the following two facts holds:
 - For all $j \in a..k$: $\mathcal{K}_{H, s_j, 0} \models \sigma_1$ and $\Box(\sigma_1 \vee \Box\sigma_2) \in \Pi'$.
 - There exists j such that $a \leq j \leq k$ and for all $h \in a..j-1$: $\mathcal{K}_{H, s_h, 0} \models \sigma_1$, and for all $h \in j..k$: $\mathcal{K}_{H, s_h, 0} \models \sigma_2$ and $\Box\sigma_2 \in \Pi'$.

Therefore, $\mathcal{K}_{H, s_a, 0} \models E\Pi$.

Let $\sigma = A\Pi \in \tau(s_a)$ be a formula of the highest-priority where $\sigma_1 \mathcal{U}(\sigma_2 \wedge \Diamond\sigma_3) \in \Pi$ is marked. By Proposition 47 and the induction hypothesis, we have that $\mathcal{K}_{H, s_v, 0} \models \sigma_2$ for some $v \geq a$, $\mathcal{K}_{H, s_z, 0} \models \sigma_1$ for all $z \in a..v-1$. In addition, $A(\Diamond\sigma_3, \Pi') \in \tau(s_v)$ (which is also the highest priority formula). Hence, by induction hypothesis, $\mathcal{K}_{H, s_v, 0} \models A(\Diamond\sigma_3, \Pi')$. Additionally, by Proposition 43 and the induction hypothesis:

- (a) For all $\sigma'_1 \mathcal{U} (\sigma'_2 \wedge \diamond \sigma'_3) \in \Pi$:
- If $\sigma'_1 \mathcal{U} (\sigma'_2 \wedge \diamond \sigma'_3)$ is also in Π' , then $\mathcal{K}_H, s_z, 0 \models \sigma'_1$ for all $z \in a..v - 1$.
 - If $\diamond \sigma'_3 \in \Pi'$ then $\mathcal{K}_H, s_z, 0 \models \sigma'_2$ for some $z \in a..v - 1$.
- (b) For all $\sigma'_1 \mathcal{U} \square \sigma'_2 \in \Pi$:
- If $\sigma'_1 \mathcal{U} \square \sigma'_2$ is also in Π' , then $\mathcal{K}_H, s_z, 0 \models \sigma'_1$ for all $z \in a..v - 1$.
 - If $\square \sigma'_2 \in \Pi'$ then there exists $j \in a..v - 1$ such that $\mathcal{K}_H, s_j, 0 \models \sigma'_1$ for all $z \in a..j - 1$ and $\mathcal{K}_H, s_z, 0 \models \sigma'_2$ for all $z \in j..v$.

Therefore, $\mathcal{K}_H, s_a, 0 \models \text{A}\Pi$.

Let $\sigma = \text{E}\Pi \in \tau(s_a)$ be a (selectable) formula of the lowest priority. If $a < i$ then, by Proposition 42 ((c) and (d)), there exists non-empty $\Pi' \in \text{Variants}(\Pi)$ such that $\text{E}\Pi' \in \tau(n_i) \subseteq \tau(s_i)$ and

- For all $\square(\sigma_1 \mathcal{U} \sigma_2) \in \Pi$: $\square(\sigma_1 \mathcal{U} \sigma_2) \in \Pi'$ and for all $z \in a..i - 1$: $\sigma_1 \in \tau(s_z)$ or $\sigma_2 \in \tau(s_z)$.
- For all $\square(\sigma_1 \vee \square \sigma_2) \in \Pi$: either $\square(\sigma_1 \vee \square \sigma_2) \in \Pi'$ and $\sigma_1 \in \tau(s_z)$ for all $z \in a..i - 1$; or $\square \sigma_2 \in \Pi'$ and there exists j such that $a \leq j \leq i$, so that for all $h \in a..j - 1$: $\sigma_1 \in \tau(s_h)$ and for all $h \in j..i$: $\sigma_2 \in \tau(s_h)$.

Since $\text{E}\Pi' \in \tau(n_1)$ is selected at some node n_h for some $i \leq h \leq j$, then by Proposition 44(c), $\sigma_2 \in \tau(s_w)$ for some $w \in i..j$. Moreover, by Proposition 42(c), for all $z \in i..j$: $\sigma_1 \in \tau(s_z)$ or $\sigma_2 \in \tau(s_z)$. In addition $\tau(n_h) = \Sigma_\ell$ for all $h \in i..j + 1$, henceforth by Proposition 42(d), for all $\square(\sigma_1 \vee \square \sigma_2) \in \Pi'$: $\sigma_1 \in \tau(s_z)$ for all $z \in i..j$. Therefore, the application of the induction hypothesis to every σ_1 and every σ_2 , allows us to ensure that $\mathcal{K}_H, s_a, 0 \models \text{E}\Pi$. The case $a \geq i$ can be seen as the particular case where $\Pi' = \Pi$ and $\text{E}\Pi \in \tau(n_h)$ for all $h \in i..j + 1$.

Let $\sigma = \text{A}\Pi \in \tau(s_a)$ be a (selectable) formula of the lowest priority. Then, Π contains at least one $\sigma_1 \mathcal{U} \square \sigma_2$ or $\square(\sigma_1 \mathcal{U} \sigma_2)$. We study two cases depending on whether there exists $\Pi' \in \text{Variants}(\Pi)$ such that $\text{A}\Pi' \in \Sigma_\ell$ or not. In the negative case, by Proposition 43 and the induction hypothesis, there exists $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond \sigma_3) \in \Pi$ and some j, j' such that $a \leq j \leq j' \leq k$, for all $h \in i..j - 1$: $\mathcal{K}_H, s_h, 0 \models \sigma_1$, $\mathcal{K}_H, s_j, 0 \models \sigma_2$, and $\mathcal{K}_H, s_{j'}, 0 \models \sigma_3$. Therefore, $\mathcal{K}_H, s_a, 0 \models \text{A}\Pi$. Otherwise, let $\text{A}\Pi' \in \Sigma_\ell \subseteq \tau(s_i)$ such that $\Pi' \in \text{Variants}(\Pi)$. By induction hypothesis and Proposition 43, $\mathcal{K}_H, s_i, 0 \models \text{A}\Pi'$ suffices to ensure that $\mathcal{K}_H, s_a, 0 \models \text{A}\Pi$. Hence, we are going to prove that $\mathcal{K}_H, s_i, 0 \models \text{A}\Pi'$. If $\text{A}\Pi'$ is non-selectable, then $\mathcal{K}_H, s_i, 0 \models \text{A}\Pi'$ holds because $\mathcal{K}_H, s_h, 0 \models \sigma_1$ for all $\square(\sigma_1 \vee \square \sigma_2) \in \Pi'$ and all $h \in i..j$. Otherwise, $\text{A}\Pi'$ is the lowest priority formula that is in the label of all the stages s_i, s_{i+1}, \dots, s_j , hence for all $h \in i..j$:

- $\sigma_1 \in \tau(s_h)$ for all $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond \sigma_3) \in \Pi'$.
- $\{\sigma_1, \sigma_2\} \subseteq \tau(s_h)$ for all $\sigma_1 \mathcal{U} \square \sigma_2 \in \Pi'$.
- $\sigma_1 \in \tau(s_h)$ or $\sigma_2 \in \tau(s_h)$ for all $\square(\sigma_1 \mathcal{U} \sigma_2) \in \Pi'$.
- $\sigma_1 \in \tau(s_h)$ for all $\square(\sigma_1 \vee \square \sigma_2) \in \Pi'$.

Moreover, $\text{A}\Pi'$ contains at least one $\sigma_1 \mathcal{U} \square \sigma_2$ or $\square(\sigma_1 \mathcal{U} \sigma_2)$, and one of its eventualities (that could be also of the form $\sigma_1 \mathcal{U} (\sigma_2 \wedge \diamond \sigma_3)$) is selected at some of the stages $s_{i_1}, s_{i_2}, \dots, s_{i_z}$. Hence, Proposition 45, along with the induction hypothesis, ensure that $\mathcal{K}_H, s_i, 0 \models \text{A}\Pi'$. \square

Corollary 50. For any expanded bunch H of $\mathcal{A}_{\Sigma_0}^{\text{sys}}$, $\mathcal{K}_H \models \Sigma_0$.

Proof. Immediate consequence of Lemma 49. \square

Now, we prove the refutational completeness of the tableau method.

Theorem 51 (Refutational completeness). For any set of state formulae Σ_0 , if $\text{UnSat}(\Sigma_0)$ then there exists a closed tableau for Σ_0 .

Proof. Suppose the contrary, that there exists no closed tableau for Σ_0 . Then the systematic tableau $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ would be open and there would be at least one expanded bunch H in $\mathcal{A}_{\Sigma_0}^{\text{sys}}$. By Corollary 50, $\mathcal{K}_B \models \Sigma_0$. Consequently Σ_0 would be satisfiable. \square

6.4. Termination

Most tableau systems for modal and temporal logics, satisfy the *analytic super-formula property (ASP)*: for every finite set of formulae Σ , there exists a finite set that contains all the formulae that may occur in any tableau for Σ . Such a set is usually called the closure of Σ . The ASP also ensures the non-existence of infinite branches where all the nodes have different labels. Hence, by controlling loops, the finiteness of proof search can be ensured. In our case, as a consequence of the β^+ -rules, the tableau system fails to satisfy the ASP, but it satisfies a slightly weaker variant which ensures completeness and that we call the *weak analytic superformula property (WASP)*: for every finite set of state formulae Σ_0 there exists a finite set (usually called the local closure of Σ) that contains all the formulae that may occur in any (systematic) tableau for Σ constructed by Algorithm \mathcal{A}^{sys} . For this purpose, the eventuality selection policy used in the \mathcal{A}^{sys} is crucial.

Theorem 52 (Termination of the tableau method). For any set of state formulae Σ_0 , the construction of the expanded tableau $\mathcal{A}_{\Sigma_0}^{\text{sys}}$ terminates.

Proof. Tableau rules produce a finite branching, hence König’s Lemma, applies. The subsumption-based simplification rules (Subsection 3.5) do prevent the generation of formulae containing the original eventuality when a “new variant” has been generated. By Propositions 44 and 45, the application of a β^+ -rule to a selected formula stops after a finite number of steps. Finally, Proposition 48 ensures that any open branch is eventually-covered. \square

Theorem 53 (Completeness of the tableau method). *For any set of state formulae Σ_0 , if Σ_0 is satisfiable then there exists a (finite) open expanded tableau for Σ_0 .*

Proof. The existence of the systematic tableau $\mathcal{A}_{\Sigma_0}^{sys}$ suffices to prove this fact, by Theorem 52. \square

7. Conclusion

We introduced a new logic, ECTL[#], in the family of BTL and its tree-style one pass tableau. This extends the expressiveness of fairness by a new class of fairness constraints with the \mathcal{U} operator. The tableau method handles inputs in an ‘analytic’ way, due to the new, crucial for branching structures, concept of ‘inner context’, in which eventualities are to be fulfilled. The tableau rules that invoke the inner context, are essential to handle A-disjunctive formulae. Our analysis of A-disjunctive and E-conjunctive formulae and of the prioritisation of eventualities, based on their structure and the context for their fulfilment, are important from the methodological point of view.

Our tableau technique is not directly extensible to CTL*. Without any significant modifications, β^+ -rules become unsound for inputs that are beyond ECTL[#] syntax due to nested path subformulae as in $A\Diamond(\circ p \wedge E\circ\neg p)$ mentioned in Fig. 1. However, for the proof of correctness of β^+ -rules, we developed the technique to identify relevant state-formulae inside the specific path-modalities. This technique will be useful in studying more expressive logics (e.g. CTL*), as it allows to identify those subformulae that do not affect the ‘context’, thus enabling the simplification of the structures.

We note that the size of the systematic tableau for the input of size m is bounded by $2^{2^{O(m^2)}}$ (see technical report at <http://www.sc.ehu.es/jiwlucaf/TechReport18.pdf>). However, the method aims at the ‘shortest’ way to fulfil the eventualities and, for many examples, finds the first open bunch, giving us a model for the tableau input. This significantly reduces the complexity. Finally, the presented technique is amenable for implementation – and this will be another stream of our future work. In the refinement and implementation of the algorithm we will be able to rely on similar techniques used in the implementation of its linear-time analogue.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We would like to thank the anonymous reviewers for their helpful comments and suggestions.

References

- [1] E.A. Emerson, J.Y. Halpern, Decision procedures and expressiveness in the temporal logic of branching time, *J. Comput. Syst. Sci.* 30 (1) (1985) 1–24.
- [2] E.A. Emerson, J.Y. Halpern, Sometimes and not never revisited: on branching versus linear time temporal logic, *J. ACM* 33 (1) (1986) 151–178.
- [3] E.M. Clarke, E.A. Emerson, A.P. Sistla, Automatic verification of finite-state concurrent systems using temporal logic specifications, *ACM Trans. Program. Lang. Syst.* 8 (2) (1986) 244–263.
- [4] B. Josko, Model checking of ctl formulae under liveness assumptions, in: T. Ottmann (Ed.), *Automata, Languages and Programming*, 14th International Colloquium, Springer-Verlag Berlin Heidelberg, Karlsruhe, 1987, pp. 5–24, Federal Republic of Germany.
- [5] E.A. Emerson, C.-L. Lei, Temporal reasoning under generalized fairness constraints, in: B. Monien, G. Vidal-Naquet (Eds.), *STACS 1986*, in: *Lecture Notes in Computer Science*, vol. 210, Springer-Verlag Berlin Heidelberg, Karlsruhe, 1986, pp. 21–37, Federal Republic of Germany.
- [6] T. Berg, H. Raffelt, Model checking, in: M. Broy, B. Jonsson, J.-P. Katoen, M. Leucker, A. Pretschner (Eds.), *Model-Based Testing of Reactive Systems*, Springer-Verlag, Berlin Heidelberg, 2005, pp. 557–603.
- [7] J. Sun, Y. Liu, J.S. Dong, H.H. Wang, Specifying and verifying event-based fairness enhanced systems, in: S. Liu, T. Maibaum, K. Araki (Eds.), *Formal Methods and Software Engineering: 10th International Conference on Formal Engineering Methods ICFEM 2008*, Springer Science and Business Media, Kitakyushu-City, Japan, 2008, pp. 5–24.
- [8] J. Kretinsky, R. Ledesma Garza, Rabinizer 2: Small deterministic automata for LTL\GU, in: *Automated Technology for Verification and Analysis – 11th International Symposium, ATVA 2013*, Springer, Heidelberg Dordrecht London New York, 2013, pp. 446–450.
- [9] E.A. Emerson, Temporal and modal logic, in: J. van Leeuwen (Ed.), *Handbook of Theoretical Computer Science* (Vol. B), MIT Press, Cambridge, USA, 1990, pp. 995–1072.
- [10] N. Markey, Temporal logics, course notes, Master Parisien de Recherche en Informatique, Paris, France, <http://www.lsv.ens-cachan.fr/Publis/PAPERS/PDF/NM-coursTL13.pdf>, 2013.
- [11] K. Brännler, M. Lange, Cut-free sequent systems for temporal logic, *J. Log. Algebraic Program.* 76 (2) (2008) 216–225.
- [12] J. Gaintzarain, M. Hermo, P. Lucio, M. Navarro, F. Orejas, Dual systems of tableaux and sequents for PLTL, *J. Log. Algebraic Program.* 78 (8) (2009) 701–722.
- [13] P. Abate, R. Goré, F. Widmann, One-pass tableaux for computation tree logic, in: N. Dershowitz, A. Voronkov (Eds.), *Logic for Programming, Artificial Intelligence, and Reasoning*, Springer Berlin Heidelberg, Berlin, Heidelberg, 2007, pp. 32–46.

- [14] O. Friedmann, M. Latte, M. Lange, A decision procedure for CTL* based on tableaux and automata, in: J. Giesl, R. Hähnle (Eds.), *Automated Reasoning*, Springer Berlin Heidelberg, Berlin, Heidelberg, 2010, pp. 331–345.
- [15] A. Pnueli, Y. Kesten, A deductive proof system for CTL*, in: L. Brim, M. Křetínský, A. Kučera, P. Jančar (Eds.), *CONCUR 2002 — Concurrency Theory*, Springer Berlin Heidelberg, Berlin, Heidelberg, 2002, pp. 24–40.
- [16] J.C. MaCabe-Dansted, M. Reynolds, Rewrite rules for CTL*, *J. Appl. Log.* 21 (2017) 24–56.
- [17] M. Reynolds, A tableau for CTL*, in: A. Cavalcanti, D. Dams (Eds.), *Proceedings. FM 2009: Formal Methods, Second World Congress*, Eindhoven, The Netherlands, November 2–6, 2009, in: *Lecture Notes in Computer Science*, vol. 5850, Springer, 2009, pp. 403–418.
- [18] M. Reynolds, A tableau-based decision procedure for CTL*, *Form. Asp. Comput.* 23 (6) (2011) 739–779.
- [19] S. Cerrito, A. David, V. Goranko, Optimal Tableau Method for Constructive Satisfiability Testing and Model Synthesis in the Alternating-Time Temporal Logic ATL+, vol. 17, 2014.
- [20] R. Gore, Tableau methods for modal and temporal logics, in: M. D’Agostino, D.M. Dov Gabbay, R. Hähnle, J. Posegga (Eds.), *Handbook of Tableau Methods*, Springer, Netherlands, Dordrecht, 1999, pp. 297–396.
- [21] A. Bolotov, M. Hermo, P. Lucio, Extending fairness expressibility of ECTL+: a tree-style one-pass tableau approach, in: N. Alechina, K. Nørvåg, W. Penczek (Eds.), *25th International Symposium on Temporal Representation and Reasoning (TIME 2018)*, in: *Leibniz International Proceedings in Informatics (LIPIcs)*, vol. 120, Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2018, 5.
- [22] R.S. Streett, E.A. Emerson, The propositional mu-calculus is elementary, in: J. Paredaens (Ed.), *Automata, Languages and Programming. Proceedings, 11th Colloquium*, Antwerp, Belgium, July 16–20, 1984, in: *Lecture Notes in Computer Science*, vol. 172, Springer, 1984, pp. 465–472.
- [23] O. Kupferman, M.Y. Vardi, P. Wolper, An automata-theoretic approach to branching-time model checking, *J. ACM* 47 (2) (2000) 312–360.